arXiv:1411.2881v1 [math.GM] 4 Nov 2014

Degenerate 4-Dimensional Matrices with Semi-Group Structure

Olga Veko*, Elena Ovsiyuk†, Alexandru Oana‡, Mircea Neagu§,
Vladimir Balan¶, Victor Red'kov‖

Abstract

While dealing with the nontrivial task of classifying Mueller matrices, of special interest is the study of the degenerate Mueller matrices (matrices with vanishing determinant, for which the law of multiplication holds, but there exists no inverse elements). Earlier, it was developed a special technique of parameterizing arbitrary 4-dimensional matrices with the use of a four 4-dimensional vector (k, m, l, n). In the following, a classification of degenerate 4-dimensional real matrices of rank 1, 2, and 3 is elaborated. To separate possible classes of degenerate matrices of ranks 1 and 2, we impose linear restrictions on (k, m, l, n), which are compatible with the group multiplication law. All the subsets of matrices obtained by this method, form either subgroups or semi-groups. To obtain singular matrices of rank 3, we specify 16 independent possibilities to get the 4-dimensional matrices with zero determinant

MSC2010: 15A66, 78A25, 35Q60, 78A99, 81V10.

Key-words: spinors; Mueller formalism; polarization optics; symmetry.

1 introduction

In polarization optics, an important issue is to classify possible classes of the Mueller matrices. In particular, of special interest are degenerate Mueller matrices (with vanishing determinant).

There is known a special technique of parameterizing arbitrary 4-dimensional matrices from $GL(4, \mathbb{C})$ with the use of four 4-dimensional vector (k, m, l, n). In the following, a classification of degenerate 4-dimensional real matrices of rank 1, 2, 3 is elaborated. To separate possible classes of degenerate matrices of ranks 1 and 2, we impose linear restrictions on (k, m, l, n), which are compatible with the group multiplication law. All the subsets of matrices obtained by this method form either subgroups or semigroups. To obtain singular matrices of rank 3, we specify 16 independent possibilities to get the 4-dimensional matrices with zero determinant[1].

In Weyl spinor basis, an arbitrary (4×4)-matrix can be parameterized by four 4-dimensional vectors (k, m, l, n)

$$G = \begin{pmatrix} k_0 + \mathbf{k}\,\vec{\sigma} & n_0 + \mathbf{n}\,\vec{\sigma} \\ l_0 + \mathbf{l}\,\vec{\sigma} & m_0 + \mathbf{m}\,\vec{\sigma} \end{pmatrix} = \begin{pmatrix} K & N \\ L & M \end{pmatrix}. \tag{1.1}$$

The matrices G will be real if the second components of parameters are imaginary

$$k_2^* = -k_2\,, \qquad m_2^* = -m_2\,, \qquad l_2^* = -l_2\,, \qquad n_2^* = -n_2\,,$$

*Kalinkovichi Gymnasium, Belarus,vekoolga@mail.ru
†Mosyr State Pedagogical University, Belarus, e.ovsiyuk@mail.ru
‡Transilvania University of Brasov, alexandru.oana@unitbv.ro
§Transilvania University of Brasov, mircea.neagu@unitbv.ro
¶University Politehnica of Bucharest, Romania, vladimir.balan@upb.ro
‖B.I. Stepanov Institute of Physics, NAS of Belarus, redkov@dragon.bas-net.by
[1]This approach is based on the works [1, 2, 3, 4, 5].

and all remaining components are real. The law of multiplication in explicit form is

$$
\begin{aligned}
k_0'' &= k_0'\, k_0 + \mathbf{k}' \cdot \mathbf{k} + n_0'\, l_0 + \mathbf{n}' \cdot \mathbf{l}\,, \\
m_0'' &= m_0'\, m_0 + \mathbf{m}' \cdot \mathbf{m} + l_0'\, n_0 + \mathbf{l}' \cdot \mathbf{n}\,, \\
n_0'' &= k_0'\, n_0 + \mathbf{k}' \cdot \mathbf{n} + n_0'\, m_0 + \mathbf{n}' \cdot \mathbf{m}\,, \\
l_0'' &= l_0'\, k_0 + \mathbf{l}' \cdot \mathbf{k} + m_0'\, l_0 + \mathbf{m}' \cdot \mathbf{l}\,,
\end{aligned}
$$

$$
\begin{aligned}
\mathbf{k}'' &= k_0'\, \mathbf{k} + \mathbf{k}'\, k_0 + i\, \mathbf{k}' \times \mathbf{k} + n_0'\, \mathbf{l} + \mathbf{n}'\, l_0 + i\, \mathbf{n}' \times \mathbf{l}\,, \\
\mathbf{m}'' &= m_0'\, \mathbf{m} + \mathbf{m}'\, m_0 + i\, \mathbf{m}' \times \mathbf{m} + l_0'\, \mathbf{n} + \mathbf{l}'\, n_0 + i\, \mathbf{l}' \times \mathbf{n}\,, \\
\mathbf{n}'' &= k_0'\, \mathbf{n} + \mathbf{k}'\, n_0 + i\, \mathbf{k}' \times \mathbf{n} + n_0'\, \mathbf{m} + \mathbf{n}'\, m_0 + i\, \mathbf{n}' \times \mathbf{m}\,, \\
\mathbf{l}'' &= l_0'\, \mathbf{k} + \mathbf{l}'\, k_0 + i\, \mathbf{l}' \times \mathbf{k} + m_0'\, \mathbf{l} + \mathbf{m}'\, l_0 + i\, \mathbf{m}' \times \mathbf{l}\,.
\end{aligned}
\tag{1.2}
$$

Now we will study the degenerate 4-dimensional matrices of rank 1, 2, 3. To separate possible classes of degenerate matrices of ranks 1 and 2, we impose linear restrictions on (k, m, l, n), which are compatible with the multiplication law (1.2). All the subsets of matrices obtained by this method, form either sub-groups or semigroups. Below, only final results will be presented.

2 One Independent Vector: Variant I(k)

First, we consider the variants with one independent vector.

Variant $I(k)$:

$$
\mathbf{n} = A\, \mathbf{k}\,,\ n_0 = \alpha\, k_0\,,\ \mathbf{m} = B\, \mathbf{k}\,,\ m_0 = \beta\, k_0\,,\ \mathbf{l} = D\, \mathbf{k}\,,\ l_0 = t\, k_0\,. \tag{2.1}
$$

The group multiplication law takes the form

$$
\begin{aligned}
k_0'' &= k_0'\, k_0 + \mathbf{k}' \cdot \mathbf{k} + \alpha t\, k_0' k_0 + AD\, \mathbf{k}' \cdot \mathbf{k}\,, \\
m_0'' &= \beta^2\, k_0' k_0 + B^2\, \mathbf{k}' \cdot \mathbf{k} + t\alpha\, k_0'\, k_0 + DA\, \mathbf{k}' \cdot \mathbf{k}\,, \\
n_0'' &= \alpha\, k_0' k_0 + A\, \mathbf{k}' \cdot \mathbf{k} + \alpha\beta\, k_0'\, k_0 + AB\, \mathbf{k}' \cdot \mathbf{k}\,, \\
l_0'' &= t\, k_0'\, k_0 + D\, \mathbf{k}' \cdot \mathbf{k} + \beta t\, k_0' k_0 + BD\, \mathbf{k}' \cdot \mathbf{k}\,,
\end{aligned}
$$

$$
\begin{aligned}
\mathbf{k}'' &= k_0'\, \mathbf{k} + \mathbf{k}'\, k_0 + i\, \mathbf{k}' \times \mathbf{k} + \alpha D\, k_0'\, \mathbf{k} + At\, \mathbf{k}' k_0 + iAD\, \mathbf{k}' \times \mathbf{k}\,, \\
\mathbf{m}'' &= \beta B\, k_0' \mathbf{k} + B\beta\, \mathbf{k}'\, k_0 + iB^2\, \mathbf{k}' \times \mathbf{k} + tA\, k_0' \mathbf{k} + D\alpha\, \mathbf{k}'\, k_0 + iDA\, \mathbf{k}' \times \mathbf{k}\,, \\
\mathbf{n}'' &= A\, k_0'\, \mathbf{k} + \alpha\, \mathbf{k}'\, k_0 + iA\, \mathbf{k}' \times \mathbf{k} + \alpha B\, k_0' \mathbf{k} + A\beta\, \mathbf{k}'\, k_0 + iAB\, \mathbf{k}' \times \mathbf{k}\,, \\
\mathbf{l}'' &= t\, k_0' \mathbf{k} + D\, \mathbf{k}' k_0 + iD\, \mathbf{k}' \times \mathbf{k} + \beta D\, k_0'\, \mathbf{k} + Bt\, \mathbf{k}'\, k_0 + iBD\, \mathbf{k}' \times \mathbf{k}\,.
\end{aligned}
$$

Let us impose the relations (2.1) for the double primed parameters:

$\underline{n_0'' = \alpha\, k_0''\,,}$

$$
\alpha(1+\beta)\, k_0' k_0 + A(1+B)\, \mathbf{k}' \cdot \mathbf{k} = \alpha(1+\alpha t)\, k_0'\, k_0 + \alpha(1+AD)\, \mathbf{k}' \cdot \mathbf{k}\,,
$$

$\underline{m_0'' = \beta\, k_0''\,,}$

$$
(\beta^2 + t\alpha)\, k_0' k_0 + (B^2 + DA)\, \mathbf{k}' \cdot \mathbf{k} = \beta(1+\alpha t)\, k_0'\, k_0 + \beta(1+AD)\, \mathbf{k}' \cdot \mathbf{k}\,,
$$

$\underline{l_0'' = t\, k_0''\,,}$

$$
t(1+\beta)\, k_0'\, k_0 + D(1+B)\, \mathbf{k}' \cdot \mathbf{k} = t\,(1+\alpha t)\, k_0'\, k_0 + t(1+AD)\, \mathbf{k}' \cdot \mathbf{k}\,,
$$

$\underline{\mathbf{n}'' = A\,\mathbf{k}''\,,}$

$$\begin{aligned}(A+\alpha B)\,k_0'\,\mathbf{k} + (\alpha + A\beta)\,\mathbf{k}'\cdot k_0 + iA(1+B)\,\mathbf{k}'\times\mathbf{k}\\ = A(1+\alpha D)\,k_0'\,\mathbf{k} + A(1+At)\,\mathbf{k}'\,k_0 + iA(1+AD)\,\mathbf{k}'\times\mathbf{k}\,,\end{aligned}$$

$\underline{\mathbf{m}'' = B\,\mathbf{k}''\,,}$

$$\begin{aligned}(\beta B + tA)\,k_0'\mathbf{k} + (B\beta + D\alpha)\,\mathbf{k}'\,k_0 + i(B^2+AD)\,\mathbf{k}'\times\mathbf{k}\\ = B(1+\alpha D)k_0'\,\mathbf{k} + B(1+At)\,\mathbf{k}'\,k_0 + iB(1+AD)\,\mathbf{k}'\times\mathbf{k}\,,\end{aligned}$$

$\underline{\mathbf{l}'' = D\,\mathbf{k}''\,,}$

$$\begin{aligned}(t+\beta D)\,k_0'\mathbf{k} + (D+Bt)\,\mathbf{k}'k_0 + iD(1+B)\,\mathbf{k}'\times\mathbf{k}\\ = D(1+\alpha D)\,k_0'\,\mathbf{k} + D(1+At)\,\mathbf{k}'\,k_0 + iD(1+AD)\,\mathbf{k}'\times\mathbf{k}\,.\end{aligned}$$

We get the equations:

$$\begin{aligned}\alpha(1+\beta) = \alpha(1+\alpha t)\,, \qquad A(1+B) = \alpha(1+AD)\,,\\ (\beta^2 + t\alpha) = \beta(1+\alpha t)\,, \qquad (B^2+DA) = \beta(1+AD)\,,\\ t(1+\beta) = t(1+\alpha t)\,, \qquad D(1+B) = t(1+AD)\,,\end{aligned}$$

$$\begin{aligned}(A+\alpha B) = A(1+\alpha D)\,,\ (\alpha + A\beta) = A(1+At)\,,\ A(1+B) = A(1+AD)\,,\\ (\beta B + tA) = B(1+\alpha D)\,,\ (B\beta + D\alpha) = B(1+At)\,,\ (B^2+AD) = B(1+AD)\,,\\ (t+\beta D) = D(1+\alpha D)\,,\ (D+Bt) = D(1+At)\,,\ D(1+B) = D(1+AD)\,.\end{aligned} \tag{2.2}$$

We are able now to find all the solutions for this system. First, let us consider the system (2.2) with three vanishing blocks – we obtain only

$\underline{\text{solution } (K-1)}$,

$$A = \alpha = 0\,,\ B = \beta = 0\,,\ D = t = 0\,, \qquad G = \begin{pmatrix} K & 0 \\ 0 & 0 \end{pmatrix}; \tag{2.3}$$

these 4-matrices form a semi-group. Their rank is equal to 2. When det $K = 0$, then the rank of G equals to 1.

Let us consider the solutions for two vanishing blocks. There arise three possibilities.

First, let it be

$$A = \alpha = 0\,, \qquad D = t = 0\,; \tag{2.4}$$

the system (2.2) gives

$$\beta^2 = \beta\,, \qquad B^2 = \beta\,, \qquad \beta B = B\,, \qquad B^2 = B\,, \tag{2.5}$$

from which we obtain one new solution when $B = \beta = 0$:

$\underline{\text{solution } (K-2)}$,

$$A = \alpha = 0\,,\ D = t = 0\,,\ D = \beta = +1\,, \qquad G = \begin{pmatrix} k_0 + \mathbf{k}\vec{\sigma} & 0 \\ 0 & k_0 + \mathbf{k}\vec{\sigma} \end{pmatrix}; \tag{2.6}$$

it is a set of non-degenerate matrices with a group structure.

Now, let it be

$$A = \alpha = 0\,, \qquad B = \beta = 0\,; \tag{2.7}$$

the system (2.2) gives

$$0 = 0\,,\ A = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ t = t\,,\ D = t\,,$$
$$0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ t = D\,,\ D = D\,,\ D = D\,,$$

so we obtain

solution $(K-3)$,

$$A = \alpha = 0\,,\ B = \beta = 0\,,\ D = t\,, G = \begin{pmatrix} K & 0 \\ DK & 0 \end{pmatrix},$$
$$G'G = \begin{pmatrix} K' & 0 \\ DK' & 0 \end{pmatrix}\begin{pmatrix} K & 0 \\ DK & 0 \end{pmatrix} = \begin{pmatrix} K'K & 0 \\ DK'K & 0 \end{pmatrix}; \tag{2.8}$$

this is a set of degenerate matrices of the rank 2 with a semi-group structure.

Now, let it be

$$B = \beta = 0\,, \qquad D = t = 0\,; \tag{2.9}$$

the system (2.2) reads

$$\alpha = \alpha\,, \qquad A = \alpha\,, \quad 0 = 0\,, \qquad 0 = 0\,, \quad 0 = 0\,, \qquad 0 = 0\,,$$
$$A = A\,,\ \alpha = A\,,\ A = A\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,\ 0 = 0\,,$$

so we get

solution $(K-4)$,

$$A = \alpha\,,\ B = \beta = 0\,,\ D = t = 0\,,\ G = \begin{pmatrix} K & AK \\ 0 & 0 \end{pmatrix},$$
$$G'G = \begin{pmatrix} K' & AK' \\ 0 & 0 \end{pmatrix}\begin{pmatrix} K & AK \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} K'K & AK'K \\ 0 & 0 \end{pmatrix}; \tag{2.10}$$

it is a set of matrices of the rank 2 and with a semi-group structure.

Let us consider cases with one vanishing block. The first possibility is

$$A = \alpha = 0\,; \tag{2.11}$$

the system (2.2) reads

$$0 = 0,\ 0 = 0,\ \beta^2 = \beta,\ B^2 = \beta,\ t(1+\beta) = t,\ D(1+B) = t,\ 0 = 0,\ 0 = 0,\ 0 = 0\,,$$
$$\beta B = B\,,\ B\beta = B\,,\ B^2 = B\,,\ (t + \beta D) = D\,,\ (D + Bt) = D\,,\ D(1+B) = D\,.$$

These equations lead to already known ones:

solution $(K-3)$,

$$A = \alpha = 0\,,\ B = \beta = 0\,,\ t = D\,,\ G = \begin{pmatrix} K & 0 \\ DK & 0 \end{pmatrix}; \tag{2.12}$$

solution $(K-2)$,

$$A = \alpha = 0\,,\ B = \beta = +1\,,\ D = t = 0\,,\ G = \begin{pmatrix} k_0 + \mathbf{k}\vec{\sigma} & 0 \\ 0 & k_0 + \mathbf{k}\vec{\sigma} \end{pmatrix}. \tag{2.13}$$

The second possibility is

$$D = t = 0 \; ; \tag{2.14}$$

the system (2.2) leads to

$$\begin{gathered} \alpha\beta = 0 \; , \; A(1+B) = \alpha \; , \; \beta^2 = \beta \; , \; B^2 = \beta \; , \\ 0 = 0 \; , \; 0 = 0 \; , \; (A + \alpha B) = A \; , \; (\alpha + A\beta) = A \; , \;\; AB = 0 \; , \\ \beta B = B \; , \; B\beta = B \; , \qquad B^2 = B \; , \; 0 = 0 \; , \; 0 = 0 \; , \; 0 = 0 \; . \end{gathered}$$

The equations (2.15) leads to already known ones:

solution $(K-2)$,

$$\begin{gathered} A = \alpha = 0 \; , \; B = +1 \; , \; \beta = +1 \; , \; D = t = 0 \; , \\ G = \begin{pmatrix} k_0 + \mathbf{k}\vec{\sigma} & 0 \\ 0 & k_0 + \mathbf{k}\vec{\sigma} \end{pmatrix} ; \end{gathered} \tag{2.15}$$

solution $(K-1)$,

$$A = \alpha \; , \; B = \beta = 0 \; , \; D = t = 0 \; , \qquad G = \begin{pmatrix} K & 0 \\ 0 & 0 \end{pmatrix} . \tag{2.16}$$

The third possibility is

$$B = \beta = 0 \; ; \tag{2.17}$$

the system (2.2) gives

$$\begin{gathered} 0 = \alpha^2 t \; , \; A = \alpha(1 + AD) \; , \; t\alpha = 0 \; , \; DA = 0 \; , \\ 0 = \alpha t^2 \; , \; D = t(1 + AD) \; , \; 0 = A\alpha D \; , \; \alpha = A(1 + At) \; , \; 0 = A^2 D \; , \\ tA = 0 \; , \; D\alpha = 0 \; , \; AD = 0 \; , \; t = D(1 + \alpha D) \; , \; 0 = DAt \; , \; 0 = AD^2 \; . \end{gathered}$$

Here there arise the already known solutions:

solution $(K-4)$,

$$A = \alpha \; , \; B = \beta = 0 \; , \; D = t = 0 \; , \qquad G = \begin{pmatrix} K & AK \\ 0 & 0 \end{pmatrix} ; \tag{2.18}$$

solution $(K-3)$,

$$A = \alpha = 0 \; , \; B = \beta = 0 \; , \; D = t \; , \qquad G = \begin{pmatrix} K & 0 \\ DK & 0 \end{pmatrix} . \tag{2.19}$$

Now, let us consider the cases in which all the blocks are non-vanishing

$$A \; , \; \alpha \neq 0 \; , \qquad B \; , \; \beta \neq 0 \; , \qquad D \; , \; t \neq 0 \; ; \tag{2.20}$$

then the system (2.2) gives

$$\begin{gathered} \beta = +\alpha t \; , \qquad A(1+B) = \alpha(1+AD) \; , \\ (\beta^2 + t\alpha) = \beta(1+\alpha t) \; , \qquad (B^2 + DA) = \beta(1+AD) \; , \\ \beta = +\alpha t \; , \qquad D(1+B) = t(1+AD) \; , \\ B = AD \; , \qquad (\alpha + A\beta) = A(1+At) \; , \qquad B = AD \; , \\ (\beta B + tA) = B(1+\alpha D) \; , \; (B\beta + D\alpha) = B(1+At) \; , \; (B^2 + AD) = B(1+AD) \; , \\ (t + \beta D) = D(1+\alpha D) \; , \qquad B = AD \; , \qquad B = AD \; . \end{gathered} \tag{2.21}$$

By eliminating B and β by $B = AD\ ,\ \beta = \alpha t$, the remaining independent equations are

$$(A-\alpha)(1+AD) = 0\ ,\ \ (AD-\alpha t)(1+AD) = 0\ ,\ \ (D-t)(1+AD) = 0\ ,$$
$$(A-\alpha)(1+At) = 0\ ,\quad (D-t)(1+\alpha D) = 0\ . \tag{2.22}$$

First, let us study the case $A = \alpha$; the system (2.22) takes the form

$$A = \alpha\ ,\qquad (D-t)(1+AD) = 0\ . \tag{2.23}$$

We get two new solutions:

solution $(K-5)$,

$$A = \alpha\ ,\ B = \beta = AD\ ,\ D = t\ ,\qquad G = \begin{pmatrix} K & AK \\ DK & ADK \end{pmatrix}; \tag{2.24}$$

it is a set of degenerate matrices of rank 4 with semi-group structure. As easily verified, the multiplication law holds indeed:

$$\begin{pmatrix} K' & AK' \\ DK' & ADK' \end{pmatrix}\begin{pmatrix} K & AK \\ DK & ADK \end{pmatrix}$$
$$= \begin{pmatrix} (K'K + ADK'K) & A(K'K + ADK'K) \\ D(K'K + ADK'K) & AD(K'K + ADK'K) \end{pmatrix}. \tag{2.25}$$

Solution $(K-5)'$,

$$A = \alpha\ ,\ B = \beta = -1\ ,\ D = t\ ,\ G = \begin{pmatrix} K & AK \\ -A^{-1}K & -K \end{pmatrix},\ G'G = 0\ ; \tag{2.26}$$

solution $(K-6)$,

$$A = \alpha\ ,\ B = -1,\ \beta = -At,\ D = -\frac{1}{A}\ ,$$
$$G = \begin{pmatrix} k_0 + \mathbf{k}\vec{\sigma} & Ak_0 + A\mathbf{k}\vec{\sigma} \\ tk_0 - A^{-1}\mathbf{k}\vec{\sigma} & Atk_0 - \mathbf{k}\vec{\sigma} \end{pmatrix}; \tag{2.27}$$

they are sets of degenerate matrices of rank 2 with semi-group structure.

Let us verify the multiplication law in the case $K-6$:

$$G'G = \begin{pmatrix} k_0' + \mathbf{k}\vec{\sigma} & Ak_0' + A\mathbf{k}'\vec{\sigma} \\ tk_0' - A^{-1}\mathbf{k}'\vec{\sigma} & Atk_0' - \mathbf{k}'\vec{\sigma} \end{pmatrix}\begin{pmatrix} k_0 + \mathbf{k}\vec{\sigma} & Ak_0 + A\mathbf{k}\vec{\sigma} \\ tk_0 - A^{-1}\mathbf{k}\vec{\sigma} & Atk_0 - \mathbf{k}\vec{\sigma} \end{pmatrix} = G''\ ;$$

the result is presented via blocks:

$$(11) = (1+At)\ k_0'k_0 + (1+At)\mathbf{k}'k_0\vec{\sigma}\ ,$$
$$(12) = A\ [(1+At)\ k_0'k_0 + (1+At)\mathbf{k}'k_0\vec{\sigma}]\ ,$$
$$(21) = t\ (1+At)\ k_0'k_0 - A^{-1}\ (1+At)\mathbf{k}'k_0\vec{\sigma}\ ,$$
$$(22) = At\ (1+At)\ k_0'k_0 - (1+At)\mathbf{k}'k_0\vec{\sigma}\ ,$$

that is

$$k_0'' = (1+At)k_0\ k_0'\ ,\qquad \mathbf{k}'' = (1+At)k_0\ \mathbf{k}'\ . \tag{2.28}$$

Let us turn back to the system (2.22) and consider the case

$$A \neq \alpha\ ,\qquad 1 + AD = 0\ ; \tag{2.29}$$

the equations (2.22) give only

solution $(K-7)$,

$$A, \alpha,\ B=-1,\ \beta=-\frac{\alpha}{A}\ ,\ D=t=-\frac{1}{A}\,,$$
$$G=\begin{pmatrix} k_0+\mathbf{k}\vec{\sigma} & \alpha k_0+A\mathbf{k}\vec{\sigma} \\ -A^{-1}(k_0+\mathbf{k}\vec{\sigma}) & -A^{-1}\alpha k_0-\mathbf{k}\vec{\sigma} \end{pmatrix}; \tag{2.30}$$

this is a set of degenerate matrices of rank 2 with semi-group structure. Let us verify the multiplication law:

$$G'G=\begin{pmatrix} k'_0+\mathbf{k}'\vec{\sigma} & \alpha k'_0+A\mathbf{k}'\vec{\sigma} \\ -A^{-1}(k'_0+\mathbf{k}'\vec{\sigma}) & -A^{-1}\alpha k'_0-\mathbf{k}'\vec{\sigma} \end{pmatrix}\begin{pmatrix} k_0+\mathbf{k}\vec{\sigma} & \alpha k_0+A\mathbf{k}\vec{\sigma} \\ -A^{-1}(k_0+\mathbf{k}\vec{\sigma}) & -A^{-1}\alpha k_0-\mathbf{k}\vec{\sigma} \end{pmatrix};$$

the result is presented by blocks

$$\begin{aligned}
(11)&=(1-\frac{\alpha}{A})\ k'_0k_0+(1-\frac{\alpha}{A})k'_0\mathbf{k}\ \vec{\sigma}\ ,\\
(12)&=\alpha\ (1-\frac{\alpha}{A})\ k'_0k_0+A\ (1-\frac{\alpha}{A})k'_0\mathbf{k}\ \vec{\sigma}\ ,\\
(21)&=-A^{-1}\ (1-\frac{\alpha}{A})\ k'_0k_0-A^{-1}\ (1-\frac{\alpha}{A})k'_0\mathbf{k}\ \vec{\sigma}\ ,\\
(22)&=-A^{-1}\alpha\ (1-\frac{\alpha}{A})\ k'_0k_0-(1-\frac{\alpha}{A})k'_0\mathbf{k}\ \vec{\sigma}\ ,
\end{aligned}$$

so we get

$$G''=G'G\ ,\qquad k''_0=(1-\frac{\alpha}{A})\ k'_0k_0\ ,\qquad \mathbf{k}''=(1-\frac{\alpha}{A})k'_0\mathbf{k}\ . \tag{2.31}$$

Thus, the analysis of the variant **I(k)** is completed.

3 One Independent Vector: Variant I(m)

Let us consider **variant** $I(m)$:

$$\mathbf{n}=A\ \mathbf{m}\ ,\ n_0=\alpha\ m_0\ ,\ \mathbf{k}=B\ \mathbf{m}\ ,\ k_0=\beta\ m_0\ ,\ \mathbf{l}=D\ \mathbf{m}\ ,\quad l_0=t\ m_0\ . \tag{3.1}$$

The group multiplication law takes the form

$$\begin{aligned}
k''_0&=\beta m'_0\ \beta m_0+B\mathbf{m}'\ B\mathbf{m}+\alpha m'_0\ tm_0+A\mathbf{m}'\ D\mathbf{m}\ ,\\
m''_0&=m'_0\ m_0+\mathbf{m}'\cdot\mathbf{m}+tm'_0\ \alpha m_0+D\mathbf{m}'\ A\mathbf{m}\ ,\\
n''_0&=\beta m'_0\ \alpha m_0+B\mathbf{m}'\ A\mathbf{m}+\alpha m'_0\ m_0+A\mathbf{m}'\cdot\mathbf{m}\ ,\\
l''_0&=tm'_0\ \beta m_0+D\mathbf{m}'\ B\mathbf{m}+m'_0\ tm_0+\mathbf{m}'\ D\mathbf{m}\ ,
\end{aligned}$$

$$\begin{aligned}
\mathbf{k}''&=\beta m'_0\ B\mathbf{m}+B\mathbf{m}'\ \beta m_0+i\ B\mathbf{m}'\times B\mathbf{m}+\alpha m'_0\ D\mathbf{m}+A\mathbf{m}'\ tm_0+i\ A\mathbf{m}'\times D\mathbf{m}\ ,\\
\mathbf{m}''&=m'_0\ \mathbf{m}+\mathbf{m}'\ m_0+i\ \mathbf{m}'\times\mathbf{m}+tm'_0\ A\mathbf{m}+D\mathbf{m}'\ \alpha m_0+i\ D\mathbf{m}'\times A\mathbf{m}\ ,\\
\mathbf{n}''&=\beta m'_0\ A\mathbf{m}+B\mathbf{m}'\ \alpha m_0+i\ B\mathbf{m}'\times A\mathbf{m}+\alpha m'_0\ \mathbf{m}+A\mathbf{m}'\ m_0+i\ A\mathbf{m}'\times\mathbf{m}\ ,\\
\mathbf{l}''&=tm'_0\ B\mathbf{m}+D\mathbf{m}'\ \beta m_0+i\ D\mathbf{m}'\times B\mathbf{m}+m'_0\ D\mathbf{m}+\mathbf{m}'\ tm_0+i\ \mathbf{m}'\times D\mathbf{m}\ .
\end{aligned}$$

We require

$$\begin{aligned}
n''_0&=\alpha\ m''_0\qquad\Longrightarrow\\
\beta m'_0\ \alpha m_0+B\mathbf{m}'\ A\mathbf{m}&+\alpha m'_0\ m_0+A\mathbf{m}'\cdot\mathbf{m}\\
=\alpha m'_0\ m_0+\alpha\mathbf{m}'\cdot\mathbf{m}&+\alpha tm'_0\ \alpha m_0+\alpha D\mathbf{m}'\ A\mathbf{m}\qquad\Longrightarrow\\
\alpha(\beta+1)=\alpha(1+\alpha t)\ ,&\qquad A(B+1)=\alpha(1+AD)\ ;
\end{aligned}$$

$$k_0'' = \beta\, m_0'' \qquad \Longrightarrow$$
$$\beta m_0'\ \beta m_0 + B\mathbf{m}'\ B\mathbf{m} + \alpha m_0'\ tm_0 + A\mathbf{m}'\ D\mathbf{m}$$
$$= \beta m_0'\ m_0 + \beta\mathbf{m}'\cdot\mathbf{m} + \beta t m_0'\ \alpha m_0 + \beta D\mathbf{m}'\ A\mathbf{m} \qquad \Longrightarrow$$
$$\beta^2 + \alpha t = \beta(1+\alpha t)\ , \qquad B^2 + AD = \beta(1+AD)\ ;$$

$$l_0'' = t\, m_0'' \qquad \Longrightarrow$$
$$tm_0'\ \beta m_0 + D\mathbf{m}'\ B\mathbf{m} + m_0'\ tm_0 + \mathbf{m}'\ D\mathbf{m}$$
$$= tm_0'\ m_0 + t\mathbf{m}'\cdot\mathbf{m} + t^2 m_0'\ \alpha m_0 + tD\mathbf{m}'\ A\mathbf{m} \qquad \Longrightarrow$$
$$t(\beta+1) = t(1+\alpha t)\ , \qquad D(B+1) = t(1+AD)\ ;$$

$$\mathbf{k}'' = B\ \mathbf{m}'' \qquad \Longrightarrow$$
$$\beta m_0'\ B\mathbf{m} + B\mathbf{m}'\ \beta m_0 + i\ B\mathbf{m}'\times B\mathbf{m} + \alpha m_0'\ D\mathbf{m} + A\mathbf{m}'\ tm_0 + i\ A\mathbf{m}'\times D\mathbf{m}$$
$$= Bm_0'\ \mathbf{m} + B\mathbf{m}'\ m_0 + i\ B\mathbf{m}'\times\mathbf{m} + Btm_0'\ A\mathbf{m} + BD\mathbf{m}'\ \alpha m_0 + i\ BD\mathbf{m}'\times A\mathbf{m}\ ,$$
$$\beta B + \alpha D = B(1+tA)\ ,\ \beta B + tA = B(1+\alpha D),\ B^2 + AD = B(1+AD)\ ;$$

$$\mathbf{n}'' = A\ \mathbf{m}'' \qquad \Longrightarrow$$
$$\beta m_0'\ A\mathbf{m} + B\mathbf{m}'\ \alpha m_0 + i\ B\mathbf{m}'\times A\mathbf{m} + \alpha m_0'\ \mathbf{m} + A\mathbf{m}'\ m_0 + i\ A\mathbf{m}'\times\mathbf{m}$$
$$= Am_0'\ \mathbf{m} + A\mathbf{m}'\ m_0 + i\ A\mathbf{m}'\times\mathbf{m} + Atm_0'\ A\mathbf{m} + AD\mathbf{m}'\ \alpha m_0 + i\ AD\mathbf{m}'\times A\mathbf{m}\ ,$$
$$\beta A + \alpha = A(1+tA)\ ,\ B\alpha + A = A(1+\alpha D)\ ,\ A(1+B) = A(1+AD)\ ;$$

$$\mathbf{l}'' = D\ \mathbf{m}'' \qquad \Longrightarrow$$
$$tm_0'\ B\mathbf{m} + D\mathbf{m}'\ \beta m_0 + i\ D\mathbf{m}'\times B\mathbf{m} + m_0'\ D\mathbf{m} + \mathbf{m}'\ tm_0 + i\ \mathbf{m}'\times D\mathbf{m}$$
$$= Dm_0'\ \mathbf{m} + D\mathbf{m}'\ m_0 + i\ D\mathbf{m}'\times\mathbf{m} + Dtm_0'\ A\mathbf{m} + DD\mathbf{m}'\ \alpha m_0 + i\ DD\mathbf{m}'\times A\mathbf{m}\ ,$$
$$tB + D = D(1+tA)\ ,\ D\beta + t = D(1+\alpha D)\ ,\ D(1+B) = D(1+AD)\ .$$

By collecting together these results, we obtain the equations

$$\begin{aligned}
\alpha(\beta+1) = \alpha(1+\alpha t)\ , &\qquad A(B+1)\alpha(1+AD)\ ,\\
= \beta^2 + \alpha t = \beta(1+\alpha t)\ , &\qquad B^2 + AD = \beta(1+AD)\ ,\\
t(\beta+1) = t(1+\alpha t)\ , &\qquad D(B+1) = t(1+AD)\ ,\\
\beta B + \alpha D = B(1+tA)\ ,\ \beta B + tA = B(1+\alpha D),&\ B^2 + AD = B(1+AD)\ ,\\
\beta A + \alpha = A(1+tA)\ ,\ B\alpha + A = A(1+\alpha D)\ ,&\ A(1+B) = A(1+AD)\ ,\\
tB + D = D(1+tA)\ ,\ D\beta + t = D(1+\alpha D)\ ,&\ D(1+B) = D(1+AD)\ .
\end{aligned} \tag{3.2}$$

The system (3.2) has a trivial solution with three vanishing blocks:

solution $(M-1)$,

$$A = \alpha = 0\ ,\ B = \beta = 0\ ,\ D = t = 0\ , \qquad G = \begin{pmatrix} 0 & 0 \\ 0 & M \end{pmatrix}. \tag{3.3}$$

Let us construct the solutions with two vanishing blocks. There are three possible variants. The first is

$$A = \alpha = 0\ , \qquad D = t = 0\ ; \tag{3.4}$$

the system (3.2) gives

$$0=0\ ,\ 0=0\ ,\ \beta^2=\beta\ ,\ B^2=\beta\ ,\ 0=0\ ,\ 0=0\ ,\ \beta B = B\ ,$$
$$\beta B = B\ ,\ B^2 = B\ ,0=0\ ,\ 0=0\ ,\ 0=0\ ,\ 0=0\ ,\ 0=0\ ,\ 0=0\ ,$$

that is

$$\beta^2 = \beta\ , \qquad B^2 = \beta\ , \qquad \beta B = B\ , \qquad B^2 = B\ . \tag{3.5}$$

The equations (3.5) have only already known solutions:

solution $(M-1)$,

$$A = \alpha = 0\ ,\ B = \beta = 0\ ,\ D = t = 0\ , \qquad G = \begin{pmatrix} 0 & 0 \\ 0 & M \end{pmatrix}; \tag{3.6}$$

solution $(M-2)$,

$$A = \alpha = 0\ ,\ B = \beta = 1\ ,\ D = t = 0\ , \qquad G = \begin{pmatrix} M & 0 \\ 0 & M \end{pmatrix}. \tag{3.7}$$

Now let us examine the variants with two vanishing blocks. The first is

$$A = \alpha = 0\ , \qquad B = \beta = 0\ ; \tag{3.8}$$

the system (3.2) gives

$$t = 0\ ,\ D = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ D = t\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ t = D\ ,\ 0 = D\ ,$$

so we get only one

solution $(M-3)$,

$$A = \alpha = 0\ ,\ B = \beta = 0\ ,\ D = t\ ,\\ G = \begin{pmatrix} 0 & 0 \\ DM & M \end{pmatrix},\ G'G = \begin{pmatrix} 0 & 0 \\ DM'M & M'M \end{pmatrix}. \tag{3.9}$$

This is a set of degenerate matrices of rank 2 with semi-group structure.

The second set is

$$D = t = 0\ , \qquad B = \beta = 0\ ; \tag{3.10}$$

the system (3.2) reads

$$0 = 0\ ,\ A = \alpha\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\\ 0 = 0\ ,\ 0 = 0,\ 0 = 0\ ,\ \alpha = A\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,$$

so we get

solution $(M-4)$,

$$A = \alpha\ ,\ B = \beta = 0\ ,\ D = t = 0\ , \qquad G = \begin{pmatrix} 0 & AM \\ 0 & M \end{pmatrix}. \tag{3.11}$$

Now let us examine the solutions with one vanishing block. There arise 3 possibilities. The first is

$$A = \alpha = 0\ ; \tag{3.12}$$

the system (3.2) gives

$$0 = 0\ ,\ 0 = 0\ ,\ \beta^2 = \beta\ ,\ B^2 = \beta\ ,\ t\beta = 0\ ,\ D(B+1) = t\ ,\ \beta B = B\ ,\ \beta B = B\ ,\\ B^2 = B\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ 0 = 0\ ,\ tB = 0\ ,\ D\beta + t = D\ ,\ DB = 0\ .$$

This system has two solutions:

solution $(M-2)$,

$$A=\alpha=0\ ,\ B=\beta=+1,\ D=t=0\ ,\qquad G=\begin{pmatrix} M & 0\\ 0 & M\end{pmatrix}; \tag{3.13}$$

solution $(M-3)$,

$$\begin{gathered} A=\alpha=0\ ,\ B=\beta=0\ ,\ D=t\ ,\\ G=\begin{pmatrix} 0 & 0\\ DM & M\end{pmatrix},\ G'G=\begin{pmatrix} 0 & 0\\ DM'M & M'M\end{pmatrix}. \end{gathered} \tag{3.14}$$

The third possibility is

$$B=\beta=0\ ; \tag{3.15}$$

the system (3.2) reads

$$\begin{gathered} 0=\alpha^2 t\ ,\ A=\alpha\ ,\ \alpha t=0\ ,\ AD=0\ ,\\ 0=\alpha t^2\ ,\ D=t\ ,\ \alpha D=0\ ,\ tA=0,\ AD=0\ ,\\ 0=tA^2\ ,\ 0=A^2D\ ,\ 0=A^2D\ ,\ 0=tDA\ ,\ 0=\alpha D^2\ ,\ 0=AD^2\ . \end{gathered}$$

These equations (3.16) have two solutions:

solution $(M-3)$,

$$\begin{gathered} A=\alpha=0\ ,\ B=\beta=0\ ,\ D=t\ ,\\ G=\begin{pmatrix} 0 & 0\\ DM & M\end{pmatrix},\ G'G=\begin{pmatrix} 0 & 0\\ DM'M & M'M\end{pmatrix}; \end{gathered} \tag{3.16}$$

solution $(M-4)$,

$$A=\alpha\ ,\ B=\beta=0\ ,\ D=t=0\ ,\ G=\begin{pmatrix} 0 & AM\\ 0 & M\end{pmatrix}. \tag{3.17}$$

Consider the last variant with one vanishing block

$$D=t=0\ ; \tag{3.18}$$

the system (3.2) gives

$$\begin{gathered} \alpha\beta=0\ ,\ A(B+1)=\alpha\ ,\ \beta^2=\beta\ ,\ B^2=\beta\ ,\\ 0=0\ ,\ 0=0\ ,\ \beta B=B\ ,\ \beta B=B,\ B^2=B\ ,\\ \beta A+\alpha=A\ ,\ B\alpha=0\ ,\ AB=0\ ,\ 0=0\ ,\ 0=0\ ,\ 0=0\ . \end{gathered} \tag{3.19}$$

These equations have two solutions:

solution $(M-2)$,

$$A=\alpha=0\ ,\ B=\beta=1\ ,\ D=t=0\ ,\ G=\begin{pmatrix} M & 0\\ 0 & M\end{pmatrix}; \tag{3.20}$$

solution $(M-4)$,

$$A=\alpha\ ,\ B=\beta=0\ ,\ D=t=0\ ,\ G=\begin{pmatrix} 0 & AM\\ 0 & M\end{pmatrix}. \tag{3.21}$$

Now, let us consider the case where all the blocks are non-vanishing

$$A\ ,\ \alpha \neq 0\ , \qquad B\ ,\ \beta \neq 0\ , \qquad D\ ,\ t \neq 0\ ; \tag{3.22}$$

The system (3.2) reads

$$\begin{aligned}
\beta = \alpha t\ , \qquad & A(B+1) = \alpha(1+AD)\ , \\
\beta^2 + \alpha t = \beta(1+\alpha t)\ , \qquad & B^2 + AD = \beta(1+AD)\ , \\
\beta = \alpha t\ , \qquad & D(B+1) = t(1+AD)\ , \\
\beta B + \alpha D = B(1+tA)\ ,\ \beta B + tA = B(1+\alpha D),\ & B^2 + AD = B(1+AD)\ , \\
\beta A + \alpha = A(1+tA)\ , \qquad & B = AD)\ , \qquad B = AD\ , \\
B = AD\ , \qquad & D\beta + t = D(1+\alpha D)\ , \qquad B = AD\ ,
\end{aligned}$$

which is equivalent to

$$\begin{aligned}
&\beta = \alpha t\ , \qquad B = AD\ ,\ (D-t)(1+AD) = 0\ , \\
&(A-\alpha)(1+AD) = 0\ ,\ (\alpha t - AD)(1+AD) = 0\ , \\
&(A-\alpha)(1+tA) = 0\ , \qquad (D-t)(1+\alpha D) = 0\ .
\end{aligned}$$

There exist two possibilities. The first is

$$\beta = \alpha t\ ,\ B = AD\ ,\ D = -\frac{1}{A}\ ,\ (A-\alpha)(1+tA) = 0\ ; \tag{3.23}$$

we get two solutions:

solution $(M-5)$,

$$\begin{aligned}
A = \alpha\ , \qquad B = -1, \qquad D = -\frac{1}{A}\ , \qquad \beta = At\ , \\
G = \begin{pmatrix} (Atm_0 - \mathbf{m}\vec{\sigma}) & (Am_0 + A\mathbf{m}\vec{\sigma}) \\ (tm_0 - A^{-1}\mathbf{m}\vec{\sigma}) & (m_0 + \mathbf{m}\vec{\sigma}) \end{pmatrix};
\end{aligned} \tag{3.24}$$

solution $(M-6)$,

$$\begin{aligned}
A\ , \alpha \neq 0\ , \qquad D = t = -\frac{1}{A}\ , \qquad B = -1\ ,\ \beta = -\frac{\alpha}{A}\ , \\
G = \begin{pmatrix} (-\alpha A^{-1} t m_0 - \mathbf{m}\vec{\sigma}) & (\alpha m_0 + A\mathbf{m}\vec{\sigma}) \\ (A^{-1}m_0 + A^{-1}\mathbf{m}\vec{\sigma}) & (m_0 + \mathbf{m}\vec{\sigma}) \end{pmatrix}.
\end{aligned} \tag{3.25}$$

The second possibility is

$$\beta = \alpha t\ , \qquad B = AD\ , \qquad D = t\ , \qquad A = \alpha\ ; \tag{3.26}$$

it leads to

solution $(M-7)$,

$$\begin{aligned}
A = \alpha \neq 0, \qquad D = t \neq 0, \qquad B = \beta = AD, \\
G = \begin{pmatrix} ADM & AM \\ DM & M \end{pmatrix},\ G'G = \begin{pmatrix} ADM' & AM' \\ DM' & M' \end{pmatrix}\begin{pmatrix} ADM & AM \\ DM & M \end{pmatrix} \\
= \begin{pmatrix} AD(1+DA)M'M & A(1+DA)M'M \\ D(1+DA)M'M & (1+DA)M'M \end{pmatrix}.
\end{aligned} \tag{3.27}$$

4 One Independent Vector: Variant I(n)

Consider **variant** $I(n)$:

$$\mathbf{k} = A\ \mathbf{n}\ ,\ k_0 = \alpha\ n_0\ ,\ \mathbf{m} = B\ \mathbf{n}\ ,\ m_0 = \beta\ n_0\ ,\ \mathbf{l} = D\ \mathbf{n}\ ,\ l_0 = t\ n_0\ . \tag{4.1}$$

The multiplication law gives

$$\begin{aligned}
k_0'' &= \alpha^2\ n_0' n_0 + A^2 \mathbf{n}' \cdot \mathbf{n} + t\ n_0' n_0 + D\ \mathbf{n}' \cdot \mathbf{n}\ , \\
m_0'' &= \beta^2\ n_0' n_0 + B^2 \mathbf{n}' \cdot \mathbf{n} + t\ n_0' n_0 + D\ \mathbf{n}' \cdot \mathbf{n}\ , \\
n_0'' &= \alpha\ n_0' n_0 + A\ \mathbf{n}' \cdot \mathbf{n} + \beta\ n_0' n_0 + B\ \mathbf{n}' \cdot \mathbf{n}\ , \\
l_0'' &= t\alpha\ n_0' n_0 + DA\ \mathbf{n}' \cdot \mathbf{n} + \beta t\ n_0' n_0 + BD\ \mathbf{n}' \cdot \mathbf{n}\ ,
\end{aligned}$$

$$\begin{aligned}
\mathbf{k}'' &= \alpha A\ n_0' \mathbf{n} + \alpha A\ \mathbf{n}' n_0 + iA^2\ \mathbf{n}' \times \mathbf{n} + D\ n_0' \mathbf{n} + t\ \mathbf{n}' n_0 + iD\ \mathbf{n}' \times \mathbf{n}\ , \\
\mathbf{m}'' &= \beta B\ n_0' \mathbf{n} + \beta B\ \mathbf{n}' n_0 + iB^2\ \mathbf{n}' \times \mathbf{n} + t\ n_0' \mathbf{n} + D\ \mathbf{n}' n_0 + iD\ \mathbf{n}' \times \mathbf{n}\ , \\
\mathbf{n}'' &= \alpha\ n_0' \mathbf{n} + A\ \mathbf{n}' n_0 + iA\ \mathbf{n}' \times \mathbf{n} + B\ n_0' \mathbf{n} + \beta\ \mathbf{n}' n_0 + iB\ \mathbf{n}' \times \mathbf{n}\ , \\
\mathbf{l}'' &= tA\ n_0' \mathbf{n} + D\alpha\ \mathbf{n}' n_0 + iDA\ \mathbf{n}' \times \mathbf{n} + \beta D\ n_0' \mathbf{n} + Bt\ \mathbf{n}' n_0 + iBD\ \mathbf{n}' \times \mathbf{n}\ .
\end{aligned}$$

We further require

$\underline{k_0'' = \alpha\ n_0'' :}$

$$\begin{aligned}
(\alpha^2 + t)\ n_0' n_0 + (A^2 + D)\ \mathbf{n}' \cdot \mathbf{n} &= \alpha(\alpha + \beta)\ n_0' n_0 + \alpha(A + B)\ \mathbf{n}' \cdot \mathbf{n}\ , \\
(\alpha^2 + t) = \alpha(\alpha + \beta)\ , \qquad (A^2 + D) &= \alpha(A + B)\ ;
\end{aligned} \tag{4.2}$$

$\underline{m_0'' = \beta\ n_0'' :}$

$$\begin{aligned}
(\beta^2 + t)\ n_0' n_0 + (B^2 + D)\ \mathbf{n}' \cdot \mathbf{n} &= \beta\ (\alpha + \beta)\ n_0' n_0 + \beta(A + B)\ \mathbf{n}' \cdot \mathbf{n}\ , \\
(\beta^2 + t) = \beta\ (\alpha + \beta)\ , \qquad (B^2 + D) &= \beta(A + B)\ ;
\end{aligned} \tag{4.3}$$

$\underline{l_0'' = t\ n_0'' :}$

$$\begin{aligned}
(t\alpha + \beta t)\ n_0' n_0 + (DA + BD)\ \mathbf{n}' \cdot \mathbf{n} &= t\ (\alpha + \beta)\ n_0' n_0 + t(A + B)\ \mathbf{n}' \cdot \mathbf{n}\ , \\
(t\alpha + \beta t) = t\ (\alpha + \beta)\ , \qquad (DA + BD) &= t(A + B)\ ;
\end{aligned} \tag{4.4}$$

$\underline{\mathbf{k}'' = A\ \mathbf{n}'' :}$

$$\begin{aligned}
&(\alpha A + D)\ n_0' \mathbf{n} + (\alpha A + t)\ \mathbf{n}' n_0 + i(A^2 + D)\ \mathbf{n}' \times \mathbf{n} \\
&= A(\alpha + B)\ n_0' \mathbf{n} + A(A + \beta)\ \mathbf{n}' n_0 + iA(A + B)\ \mathbf{n}' \times \mathbf{n}\ , \\
&(\alpha A + D) = A(\alpha + B),\ (\alpha A + t) = A(A + \beta)\ ,\ (A^2 + D) = A(A + B)\ ;
\end{aligned} \tag{4.5}$$

$\underline{\mathbf{m}'' = B\ \mathbf{n}'' :}$

$$\begin{aligned}
&(\beta B + t)\ n_0' \mathbf{n} + (\beta B + D)\ \mathbf{n}' n_0 + i(B^2 + D)\ \mathbf{n}' \times \mathbf{n} \\
&= B(\alpha + B)\ n_0' \mathbf{n} + B(A + \beta)\ \mathbf{n}' n_0 + iB(A + B)\ \mathbf{n}' \times \mathbf{n}\ ,
\end{aligned}$$

$$(\beta B + t) = B(\alpha + B)\ ,\ (\beta B + D) = B(A + \beta)\ ,\ (B^2 + D) = B(A + B)\ ; \tag{4.6}$$

$\underline{\mathbf{l}'' = D\,\mathbf{n}'' :}$

$$
\begin{aligned}
&(tA+\beta D)\,n_0'\mathbf{n} + (D\alpha + Bt)\,\mathbf{n}' n_0 + i(DA+BD)\,\mathbf{n}'\times\mathbf{n} \\
&= D(\alpha+B)\,n_0'\mathbf{n} + D(A+\beta)\,\mathbf{n}' n_0 + iD(A+B)\,\mathbf{n}'\times\mathbf{n}\,,
\end{aligned}
$$

$$
(tA+\beta D) = D(\alpha+B)\,,\ (D\alpha+Bt) = D(A+\beta)\,,\ (DA+BD) = D(A+B)\,. \tag{4.7}
$$

Thus, we have obtained the system

$$
\begin{aligned}
(\alpha^2+t) = \alpha(\alpha+\beta)\,, &\qquad (A^2+D) = \alpha(A+B)\,, \\
(\beta^2+t) = \beta\,(\alpha+\beta)\,, &\qquad (B^2+D) = \beta(A+B)\,, \\
t(\alpha+\beta) = t\,(\alpha+\beta)\,, &\qquad D(A+B) = t(A+B)\,, \\
(\alpha A+D) = A(\alpha+B)\,,\ (\alpha A+t) = A(A+\beta)\,,\ &(A^2+D) = A(A+B)\,, \\
(\beta B+t) = B(\alpha+B)\,,\ (\beta B+D) = B(A+\beta)\,,\ &(B^2+D) = B(A+B)\,, \\
(tA+\beta D) = D(\alpha+B)\,,\ (D\alpha+Bt) = D(A+\beta)\,,\ &D(A+B) = D(A+B)\,,
\end{aligned}
$$

which is equivalent to

$$
\begin{aligned}
t = \alpha\beta\,,\ (A^2+D) = \alpha(A+B)\,,\ &(B^2+D) = \beta(A+B)\,, \\
(D-t)(A+B) = 0\,,\ D = AB,\ &(\alpha A+t) = A(A+\beta)\,, \\
(\beta B+t) = B(\alpha+B)\,,\ (tA+\beta D) = D(\alpha+B)\,,\ &(D\alpha+Bt) = D(A+\beta)\,.
\end{aligned}
$$

By eliminating the variables D and t, we get

$$
D = AB\,, \qquad t = \alpha\beta\,; \tag{4.8}
$$

the remaining independent equations are

$$
\begin{aligned}
(A-\alpha)(A+B) &= 0\,, \\
(B-\beta)(B+A) &= 0\,, \\
(AB-\alpha\beta)(A+B) &= 0\,, \\
(A-\alpha)(A+\beta) &= 0\,, \\
(B-\beta)(B+\alpha) &= 0\,, \\
A(\beta-B)(\alpha+B) &= 0\,, \\
B(\alpha-A)(A+\beta) &= 0\,.
\end{aligned} \tag{4.9}
$$

First, assume that

$$
A+B \neq 0\,, \tag{4.10}
$$

then (4.9) give

$$
A = \alpha\,, \qquad B = \beta\,, \qquad B(A-B) = 0\,. \tag{4.11}
$$

Here, there arise two solutions:

$\underline{\text{solution } (N-1)}$,

$$
A = \alpha\,,\ B = \beta = 0,\ D = t = 0, \qquad G = \begin{pmatrix} AN & N \\ 0 & 0 \end{pmatrix}; \tag{4.12}
$$

solution $(N-2)$,

$$A=\alpha\ ,\ B=\beta=A,\ D=t=A^2,\qquad G=\begin{pmatrix} AN & N \\ A^2N & AN \end{pmatrix},$$
$$G'G=\begin{pmatrix} AN' & N' \\ A^2N' & AN' \end{pmatrix}\begin{pmatrix} AN & N \\ A^2N & AN \end{pmatrix}=\begin{pmatrix} A\,2AN'N & 2AN'N \\ A^2\,2AN'N & A\,2AN'N \end{pmatrix}. \tag{4.13}$$

Now, let it be

$$A+B=0\ ,\qquad B=-A\ ; \tag{4.14}$$

correspondingly to (4.9) it follows

$$\begin{aligned} B=-A\ ,\qquad D=-A^2\ ,\qquad t=\alpha\beta\ ,\\ (A-\alpha)(A+\beta)=0\ ,\\ A(\beta+A)(\alpha-A)=0\ ,\\ A(\alpha-A)(A+\beta)=0\ . \end{aligned} \tag{4.15}$$

Two solutions are possible:
solution $(N-3)$,

$$\begin{aligned} \beta=B=-A\ ,\qquad D=-A^2\ ,\qquad t=-\alpha A\ ,\\ G=\begin{pmatrix} \alpha n_0+A\mathbf{n}\vec{\sigma} & n_0+\mathbf{n}\vec{\sigma} \\ -\alpha A n_0-A^2\mathbf{n}\vec{\sigma} & -An_0-A\mathbf{n}\vec{\sigma} \end{pmatrix},\qquad \det\ G=0\ ; \end{aligned} \tag{4.16}$$

solution $(N-4)$,

$$\begin{aligned} A=+\alpha\ ,\qquad B=-A\ ,\qquad D=-A^2\ ,\qquad t=A\beta\ ,\\ G=\begin{pmatrix} An_0+A\mathbf{n}\vec{\sigma} & n_0+\mathbf{n}\vec{\sigma} \\ \beta A n_0-A^2\mathbf{n}\vec{\sigma} & \beta n_0-A\mathbf{n}\vec{\sigma} \end{pmatrix},\qquad \det\ G=0\ . \end{aligned} \tag{4.17}$$

5 One Independent Vector: Variant I(l)

Let us examine the **variant** $I(l)$:

$$\mathbf{k}=A\,\mathbf{l}\ ,\ k_0=\alpha\ l_0\ ,\ \mathbf{m}=B\,\mathbf{l}\ ,\ m_0=\beta\ l_0\ ,\ \mathbf{n}=D\,\mathbf{l}\ ,\ n_0=t\ l_0\ . \tag{5.1}$$

The multiplication law takes the form

$$\begin{aligned} k_0''=\alpha^2 l_0'\ l_0+A^2\mathbf{l}'\cdot\mathbf{l}+tl_0'\ l_0+D\mathbf{l}'\cdot\mathbf{l}\ ,\\ m_0''=\beta^2 l_0'\ l_0+B^2\mathbf{l}'\cdot\mathbf{l}+tl_0'\ l_0+D\mathbf{l}'\cdot\mathbf{l}\ ,\\ n_0''=\alpha t l_0'\ l_0+AD\mathbf{l}'\cdot\mathbf{l}+\beta t l_0'\ l_0+BD\mathbf{l}'\cdot\mathbf{l}\ ,\\ l_0''=\alpha l_0'\ l_0+A\mathbf{l}'\cdot\mathbf{l}+\beta l_0'\ l_0+B\mathbf{l}'\cdot\mathbf{l}\ , \end{aligned} \tag{5.2}$$

$$\begin{aligned} \mathbf{k}''=A\alpha l_0'\,\mathbf{l}+A\alpha\mathbf{l}'\ l_0+i\ A^2\mathbf{l}'\times\mathbf{l}+tl_0'\,\mathbf{l}+D\ \mathbf{l}'\ l_0+i\ D\mathbf{l}'\times\mathbf{l}\ ,\\ \mathbf{m}''=B\beta l_0'\,\mathbf{l}+B\beta\mathbf{l}'\ l_0+i\ B^2\mathbf{l}'\times\mathbf{l}+Dl_0'\,\mathbf{l}+t\mathbf{l}'\ l_0+i\ D\mathbf{l}'\times\mathbf{l}\ ,\\ \mathbf{n}''=\alpha Dl_0'\,\mathbf{l}+At\mathbf{l}'\ l_0+i\ AD\mathbf{l}'\times\mathbf{l}+Btl_0'\,\mathbf{l}+\beta D\mathbf{l}'\ l_0+i\ BD\mathbf{l}'\times\mathbf{l}\ ,\\ \mathbf{l}''=Al_0'\,\mathbf{l}+\alpha\mathbf{l}'\ l_0+i\ A\mathbf{l}'\times\mathbf{l}+\beta l_0'\,\mathbf{l}+B\mathbf{l}'\ l_0+i\ B\mathbf{l}'\times\mathbf{l}\ . \end{aligned} \tag{5.3}$$

We further require

$$
\begin{aligned}
k_0'' = \alpha\, l_0'' \qquad &\Longrightarrow \\
(\alpha^2+t) l_0' \, l_0 + (A^2+D)\mathbf{l}'\cdot\mathbf{l} = \alpha(\alpha+\beta) l_0' \, l_0 + \alpha(A+B)\mathbf{l}'\cdot\mathbf{l} \quad &\Longrightarrow \\
(\alpha^2+t) = \alpha(\alpha+\beta)\,, \qquad (A^2+D) = \alpha(A+B)\,; &
\end{aligned}
$$

$$
\begin{aligned}
m_0'' = \beta\, l_0'' \qquad &\Longrightarrow \\
(\beta^2+t) l_0' \, l_0 + (B^2+D)\mathbf{l}'\cdot\mathbf{l} = \beta(\alpha+\beta) l_0' \, l_0 + \beta(A+B)\mathbf{l}'\cdot\mathbf{l} \quad &\Longrightarrow \\
(\beta^2+t) = \beta(\alpha+\beta)\,, \qquad (B^2+D) = \beta(A+B)\,; &
\end{aligned}
$$

$$
\begin{aligned}
n_0'' = t\, l_0'' \qquad &\Longrightarrow \\
(\alpha t+\beta t) l_0' \, l_0 + (AD+BD)\mathbf{l}'\cdot\mathbf{l} = t(\alpha+\beta) l_0' \, l_0 + t(A+B)\mathbf{l}'\cdot\mathbf{l} \quad &\Longrightarrow \\
(\alpha t+\beta t) = t(\alpha+\beta)\,, \qquad (AD+BD) = t(A+B)\,; &
\end{aligned}
$$

$$
\begin{aligned}
\mathbf{k}'' = A\,\mathbf{l}'' \qquad &\Longrightarrow \\
(A\alpha+t) l_0' \, \mathbf{l} + (A\alpha+D)\mathbf{l}' \, l_0 + i\,(A^2+D)\mathbf{l}'\times\mathbf{l} & \\
= A(A+\beta) l_0' \, \mathbf{l} + A(\alpha+B)\mathbf{l}' \, l_0 + i\,A(A+B)\mathbf{l}'\times\mathbf{l} \quad &\Longrightarrow \\
(A\alpha+t) = A(A+\beta)\,, \qquad (A\alpha+D) = A(\alpha+B)\,, \qquad (A^2+D) = A(A+B)\,; &
\end{aligned}
$$

$$
\begin{aligned}
\mathbf{m}'' = B\,\mathbf{l}'' \qquad &\Longrightarrow \\
(B\beta+D) l_0' \, \mathbf{l} + (B\beta+t)\mathbf{l}' \, l_0 + i\,(B^2+D)\mathbf{l}'\times\mathbf{l} & \\
= B(A+\beta) l_0' \, \mathbf{l} + B(\alpha+B)\mathbf{l}' \, l_0 + i\,B(A+B)\mathbf{l}'\times\mathbf{l} \quad &\Longrightarrow \\
(B\beta+D) = B(A+\beta)\,, \qquad (B\beta+t) = B(\alpha+B)\,, \qquad (B^2+D) = B(A+B)\,; &
\end{aligned}
$$

$$
\begin{aligned}
\mathbf{n}'' = D\,\mathbf{l}'' \qquad &\Longrightarrow \\
(\alpha D+Bt) l_0' \, \mathbf{l} + (At+\beta D)\mathbf{l}' \, l_0 + i\,(AD+BD)\mathbf{l}'\times\mathbf{l} & \\
= D(A+\beta) l_0' \, \mathbf{l} + D(\alpha+B)\mathbf{l}' \, l_0 + i\,D(A+B)\mathbf{l}'\times\mathbf{l} \quad &\Longrightarrow \\
(\alpha D+Bt) = D(A+\beta)\,,\ (At+\beta D) = D(\alpha+B)\,,\ (AD+BD) = D(A+B)\,. &
\end{aligned}
$$

By collecting the results together, we get

$$
\begin{aligned}
(\alpha^2+t) = \alpha(\alpha+\beta)\,, \qquad (A^2+D) = \alpha(A+B)\,, & \\
(\beta^2+t) = \beta(\alpha+\beta)\,, \qquad (B^2+D) = \beta(A+B)\,, & \\
(\alpha t+\beta t) = t(\alpha+\beta)\,, \qquad (AD+BD) = t(A+B)\,, & \\
(A\alpha+t) = A(A+\beta)\,,\ (A\alpha+D) = A(\alpha+B)\,,\ (A^2+D) = A(A+B)\,, & \\
(B\beta+D) = B(A+\beta)\,,\ (B\beta+t) = B(\alpha+B)\,,\ (B^2+D) = B(A+B)\,, & \\
(\alpha D+Bt) = D(A+\beta)\,,\ (At+\beta D) = D(\alpha+B)\,,\ (AD+BD) = D(A+B)\,. &
\end{aligned}
\tag{5.4}
$$

The system (5.4) is equivalent to

$$
\begin{aligned}
t = \alpha\beta\,, \qquad (A^2+D) = \alpha(A+B)\,, & \\
t = \beta\alpha\,, \qquad (B^2+D) = \beta(A+B)\,, & \\
0 = 0\,, \qquad (D-t)(A+B) = 0\,, & \\
(A\alpha+t) = A(A+\beta)\,, \qquad (A\alpha+D) = A(\alpha+B)\,, \qquad D = AB\,, & \\
(B\beta+D) = B(A+\beta)\,, \qquad (B\beta+t) = B(\alpha+B)\,, \qquad D = AB\,, & \\
(\alpha D+Bt) = D(A+\beta)\,, \qquad (At+\beta D) = D(\alpha+B)\,, \qquad 0 = 0\,. &
\end{aligned}
\tag{5.5}
$$

By excluding the variables t and D via

$$t = \alpha\beta \ , \qquad D = AB \ , \tag{5.6}$$

we obtain

$$\begin{gathered}(A-\alpha)(A+B)=0 \ , \ (B-\beta)(A+B)=0 \ , \ (AB-\alpha\beta)(A+B)=0 \ , \\ (A-\alpha)(A+\beta)=0 \ , \quad (B-\beta)(\alpha+B)=0 \ , \\ A(\beta-B)(\alpha+B)=0 \ , \quad B(\alpha-A)(A+\beta)=0 \ . \end{gathered} \tag{5.7}$$

Because these equations coincide with (4.9), we may use yet the known results:

solution $(L-1)$,

$$\begin{gathered}A=\alpha \ , \ B=\beta=0 \ , \ D=t=0 \ , \qquad G=\begin{pmatrix} AL & 0 \\ L & 0 \end{pmatrix}, \\ G'G=\begin{pmatrix} AL & 0 \\ L & 0 \end{pmatrix}\begin{pmatrix} AL' & L' \\ 0 & 0 \end{pmatrix}=\begin{pmatrix} AL'L & 0 \\ L'L & 0 \end{pmatrix}; \end{gathered} \tag{5.8}$$

solution $(L-2)$,

$$\begin{gathered}A=\alpha \ , \ B=\beta=A \ , \ D=t=A^2 \ , \qquad G=\begin{pmatrix} AL & A^2L \\ L & AL \end{pmatrix}, \\ G'G=\begin{pmatrix} AL' & A^2L' \\ L' & AL' \end{pmatrix}\begin{pmatrix} AL & A^2L \\ L & AL \end{pmatrix}=\begin{pmatrix} A\,2AL'L & A^2\,2AL'L \\ 2AL'L & A\,2AL'L \end{pmatrix}; \end{gathered} \tag{5.9}$$

solution $(L-3)$,

$$\begin{gathered}\beta=B=-A \ , \qquad D=-A^2 \ , \qquad t=-\alpha A \ , \\ G=\begin{pmatrix} \alpha l_0+A\mathbf{l}\vec{\sigma} & -\alpha A l_0-A^2\mathbf{l} \\ l_0+\mathbf{l}\vec{\sigma} & -Al_0-A\mathbf{l}\vec{\sigma} \end{pmatrix}, \qquad \det\ G=0 \ ; \end{gathered} \tag{5.10}$$

solution $(L-4)$,

$$\begin{gathered}A=+\alpha \ , \qquad B=-A \ , \qquad D=-A^2 \ , \qquad t=A\beta \ , \\ G=\begin{pmatrix} Al_0+A\mathbf{l}\vec{\sigma} & \beta A l_0-A^2\mathbf{l}\vec{\sigma} \\ l_0+\mathbf{l}\vec{\sigma} & \beta l_0-A\mathbf{l}\vec{\sigma} \end{pmatrix}, \qquad \det\ G=0 \ . \end{gathered} \tag{5.11}$$

6 Two Independent Vectors: Variant II(k,m)

Let us examine the **variant II(k,m)**:

$$\begin{aligned}\mathbf{n}&=A\mathbf{k}+B\mathbf{m} \ , \qquad & n_0&=\alpha k_0+\beta m_0 \ , \\ \mathbf{l}&=C\mathbf{k}+D\mathbf{m} \ , \qquad & l_0&=sk_0+tm_0 \ . \end{aligned} \tag{6.1}$$

The multiplication law takes the form

$$\begin{aligned}k_0''&=k_0'\,k_0+\mathbf{k}'\cdot\mathbf{k}+(\alpha k_0'+\beta m_0')(sk_0+tm_0)+(A\mathbf{k}'+B\mathbf{m}')(C\mathbf{k}+D\mathbf{m}) \ , \\ m_0''&=m_0'\,m_0+\mathbf{m}'\cdot\mathbf{m}+(sk_0'+tm_0')(\alpha k_0+\beta m_0+(C\mathbf{k}'+D\mathbf{m}')(A\mathbf{k}+B\mathbf{m}) \ , \\ n_0''&=k_0'(\alpha k_0+\beta m_0)+\mathbf{k}'(A\mathbf{k}+B\mathbf{m})+(\alpha k_0'+\beta m_0')m_0+(A\mathbf{k}'+B\mathbf{m}')\mathbf{m} \ , \\ l_0''&=(sk_0'+tm_0')k_0+(C\mathbf{k}'+D\mathbf{m}')\mathbf{k}+m_0'(sk_0+tm_0)+\mathbf{m}'(C\mathbf{k}+D\mathbf{m}) \ , \end{aligned}$$

$$
\begin{aligned}
\mathbf{k}'' &= k_0'\,\mathbf{k} + \mathbf{k}'\,k_0 + i\,\mathbf{k}'\times\mathbf{k} + (\alpha k_0' + \beta m_0')(C\mathbf{k} + D\mathbf{m}) \\
&+(A\mathbf{k}' + B\mathbf{m}')(sk_0 + tm_0) + i\,(A\mathbf{k}' + B\mathbf{m}')\times(C\mathbf{k} + D\mathbf{m})\,, \\
\mathbf{m}'' &= m_0'\,\mathbf{m} + \mathbf{m}'\,m_0 + i\,\mathbf{m}'\times\mathbf{m} + (sk_0' + tm_0')(A\mathbf{k} + B\mathbf{m}) \\
&+(C\mathbf{k}' + D\mathbf{m}')(\alpha k_0 + \beta m_0) + i\,(C\mathbf{k}' + D\mathbf{m}')\times(A\mathbf{k} + B\mathbf{m})\,, \\
\mathbf{n}'' &= k_0'(A\mathbf{k} + B\mathbf{m}) + \mathbf{k}'(\alpha k_0 + \beta m_0) + i\,\mathbf{k}'\times(A\mathbf{k} + B\mathbf{m}) \\
&+(\alpha k_0' + \beta m_0')\mathbf{m} + (A\mathbf{k}' + B\mathbf{m}')m_0 + i\,(A\mathbf{k}' + B\mathbf{m}')\times\mathbf{m}\,, \\
\mathbf{l}'' &= (sk_0' + tm_0')\mathbf{k} + (C\mathbf{k}' + D\mathbf{m}')k_0 + i\,(C\mathbf{k}' + D\mathbf{m}')\times\mathbf{k} \\
&+m_0'(C\mathbf{k} + D\mathbf{m}) + \mathbf{m}'(sk_0 + tm_0) + i\,\mathbf{m}'\times(C\mathbf{k} + D\mathbf{m})\,.
\end{aligned}
$$

By requiring $\alpha k_0'' + \beta m_0'' = n_0''$, or

$$
\begin{aligned}
&\alpha\,[\,k_0'\,k_0 + \mathbf{k}'\cdot\mathbf{k} + (\alpha k_0' + \beta m_0')(sk_0 + tm_0) + (A\mathbf{k}' + B\mathbf{m}')(C\mathbf{k} + D\mathbf{m})\,] \\
&+\beta\,[\,m_0'\,m_0 + \mathbf{m}'\cdot\mathbf{m} + (sk_0' + tm_0')(\alpha k_0 + \beta m_0 + (C\mathbf{k}' + D\mathbf{m}')(A\mathbf{k} + B\mathbf{m})\,] \\
&= k_0'(\alpha k_0 + \beta m_0) + \mathbf{k}'(A\mathbf{k} + B\mathbf{m}) + (\alpha k_0' + \beta m_0')m_0 + (A\mathbf{k}' + B\mathbf{m}')\mathbf{m}\,,
\end{aligned}
$$

we obtain

$$
\begin{array}{llll}
k_0'k_0: & \alpha + \alpha^2 s + \alpha\beta s = \alpha\,, & m_0'm_0: & \alpha\beta t + \beta + t\beta^2 = \beta\,, \\
k_0'm_0: & \alpha^2 t + \beta^2 s = \beta + \alpha\,, & m_0'k_0: & \alpha\beta s + \alpha\beta t = 0\,, \\
\mathbf{m}'\cdot\mathbf{k}: & \alpha BC + \beta AD = 0\,, & \mathbf{k}'\cdot\mathbf{m}: & \alpha AD + \beta CB = B + A\,, \\
\mathbf{k}'\cdot\mathbf{k}: & \alpha + \alpha AC + \beta AC = A\,, & \mathbf{m}'\cdot\mathbf{m}: & \alpha BD + \beta + \beta DB = B\,.
\end{array}
\tag{6.2}
$$

We further require $sk_0'' + tm_0'' = l_0''$, or

$$
\begin{aligned}
&s\,[\,k_0'\,k_0 + \mathbf{k}'\cdot\mathbf{k} + (\alpha k_0' + \beta m_0')(sk_0 + tm_0) + (A\mathbf{k}' + B\mathbf{m}')(C\mathbf{k} + D\mathbf{m})\,] \\
&+t\,[\,m_0'\,m_0 + \mathbf{m}'\cdot\mathbf{m} + (sk_0' + tm_0')(\alpha k_0 + \beta m_0 + (C\mathbf{k}' + D\mathbf{m}')(A\mathbf{k} + B\mathbf{m})\,] \\
&= (sk_0' + tm_0')k_0 + (C\mathbf{k}' + D\mathbf{m}')\mathbf{k} + m_0'(sk_0 + tm_0) + \mathbf{m}'(C\mathbf{k} + D\mathbf{m})\,,
\end{aligned}
$$

and we find

$$
\begin{array}{llll}
k_0'k_0: & s + s^2\alpha + ts\alpha = s\,, & m_0'm_0: & \beta st + t + \beta t^2 = t\,, \\
k_0'm_0: & st\alpha + st\beta = 0\,, & m_0'k_0: & \beta s^2 + \alpha t^2 = t + s\,, \\
\mathbf{k}'\cdot\mathbf{m}: & sAD + tCB = 0, & \mathbf{m}'\cdot\mathbf{k}: & sBC + tAD = C + D\,, \\
\mathbf{k}'\cdot\mathbf{k}: & s + sAC + tAC = C\,, & \mathbf{m}'\cdot\mathbf{m}: & sBD + t + tBD = D\,.
\end{array}
\tag{6.3}
$$

By requiring $A\mathbf{k}'' + B\mathbf{m}'' = \mathbf{n}''$, or

$$
\begin{aligned}
&A\,[\,k_0'\,\mathbf{k} + \mathbf{k}'\,k_0 + i\,\mathbf{k}'\times\mathbf{k} + (\alpha k_0' + \beta m_0')(C\mathbf{k} + D\mathbf{m}) \\
&+(A\mathbf{k}' + B\mathbf{m}')(sk_0 + tm_0) + i\,(A\mathbf{k}' + B\mathbf{m}')\times(C\mathbf{k} + D\mathbf{m})\,] \\
&+B\,[\,m_0'\,\mathbf{m} + \mathbf{m}'\,m_0 + i\,\mathbf{m}'\times\mathbf{m} + (sk_0' + tm_0')(A\mathbf{k} + B\mathbf{m}) \\
&+(C\mathbf{k}' + D\mathbf{m}')(\alpha k_0 + \beta m_0) + i\,(C\mathbf{k}' + D\mathbf{m}')\times(A\mathbf{k} + B\mathbf{m})\,] \\
&= k_0'(A\mathbf{k} + B\mathbf{m}) + \mathbf{k}'(\alpha k_0 + \beta m_0) + i\,\mathbf{k}'\times(A\mathbf{k} + B\mathbf{m}) \\
&+(\alpha k_0' + \beta m_0')\mathbf{m} + (A\mathbf{k}' + B\mathbf{m}')m_0 + i\,(A\mathbf{k}' + B\mathbf{m}')\times\mathbf{m}\,,
\end{aligned}
$$

we yield

$$
\begin{array}{llll}
k_0'\mathbf{k}: & A + \alpha AC + sBA = A\,, & \mathbf{m}'m_0: & tAB + B + \beta BD = B\,, \\
\mathbf{k}'k_0: & A + sA^2 + \alpha BC = \alpha\,, & m_0'\mathbf{m}: & \beta AD + B + tB^2 = \beta\,, \\
k_0'\mathbf{m}: & \alpha AD + sB^2 = B + \alpha\,, & \mathbf{k}'m_0: & tA^2 + \beta CB = \beta + A\,, \\
m_0'\mathbf{k}: & \beta AC + tAB = 0\,, & +\mathbf{m}'k_0: & sAB + \alpha BD = 0\,, \\
\mathbf{k}'\times\mathbf{k}: & A + A^2C + BCA = A\,, & \mathbf{m}'\times\mathbf{m}: & ABD + B + DB^2 = B\,, \\
\mathbf{k}'\times\mathbf{m}: & A^2D + CB^2 = B + A\,, & \mathbf{m}'\times\mathbf{k}: & ABC + ADB = 0\,.
\end{array}
$$

Now, requiring $C\mathbf{k}'' + D\mathbf{m}'' = \mathbf{l}''$, or

$$\begin{aligned}
&C\,[\;k_0'\;\mathbf{k}+\mathbf{k}'\;k_0+i\;\mathbf{k}'\times\mathbf{k}+(\alpha k_0'+\beta m_0')(C\mathbf{k}+D\mathbf{m})\\
&+(A\mathbf{k}'+B\mathbf{m}')(sk_0+tm_0)+i\;(A\mathbf{k}'+B\mathbf{m}')\times(C\mathbf{k}+D\mathbf{m})]\\
&+D\,[\;m_0'\;\mathbf{m}+\mathbf{m}'\;m_0+i\;\mathbf{m}'\times\mathbf{m}+(sk_0'+tm_0')(A\mathbf{k}+B\mathbf{m})\\
&+(C\mathbf{k}'+D\mathbf{m}')(\alpha k_0+\beta m_0)+i\;(C\mathbf{k}'+D\mathbf{m}')\times(A\mathbf{k}+B\mathbf{m})\\
&=(sk_0'+tm_0')\mathbf{k}+(C\mathbf{k}'+D\mathbf{m}')k_0+i\;(C\mathbf{k}'+D\mathbf{m}')\times\mathbf{k}\\
&+m_0'(C\mathbf{k}+D\mathbf{m})+\mathbf{m}'(sk_0+tm_0)+i\;\mathbf{m}'\times(C\mathbf{k}+D\mathbf{m})\;,
\end{aligned}$$

we find

$$\begin{aligned}
k_0'\mathbf{k}:&\quad C+\alpha C^2+sAD=s\;,\quad \mathbf{m}'m_0:\quad tCB+D+\beta D^2=t\;,\\
\mathbf{k}'k_0:&\quad C+sAC+\alpha CD=C\;,\quad m_0'\mathbf{m}:\quad \beta CD+D+tBD=D\;,\\
k_0'\mathbf{m}:&\quad \alpha DC+sBD=0\;,\quad \mathbf{k}'m_0:\quad tAC+\beta CD=0\;,\\
m_0'\mathbf{k}:&\quad \beta C^2+tAD=t+C\;,\quad +\mathbf{m}'k_0:\quad sCB+\alpha D^2=D+s\;,\\
\mathbf{k}'\times\mathbf{k}:&\quad C+AC^2+DCA=C\;,\quad \mathbf{m}'\times\mathbf{m}:\quad CBD+D+BD^2=D\;,\\
\mathbf{k}'\times\mathbf{m}:&\quad ACD+CBD=0\;,\quad \mathbf{m}'\times\mathbf{k}:\quad BC^2+AD^2=D+C\;.
\end{aligned}$$

We collect the results together:

$$\begin{aligned}
&\alpha s(\alpha+\beta)=0\;, && \beta t(\alpha+\beta)=0\;,\\
&\alpha^2t+\beta^2s=\beta+\alpha\;, && \alpha\beta(s+t)=0\;,\\
&\alpha BC+\beta AD=0\;, && \alpha AD+\beta BC=B+A\;,\\
&\alpha+(\alpha+\beta)AC=A\;, && \beta+(\alpha+\beta)BD=B\;,\\
&\\
&s\alpha(s+t)=0\;, && \beta t(s+t)=0\;,\\
&st(\alpha+\beta)=0\;, && \beta s^2+\alpha t^2=t+s\;,\\
&sAD+tCB=0, && sBC+tAD=C+D\;,\\
&s+AC(s+t)=C\;, && t+BD(t+s)=D\;,\\
&\\
&A(\alpha C+sB)=0\;, && B(\beta D+tA)=0\;,\\
&A+sA^2+\alpha BC=\alpha\;, && B+tB^2+\beta AD=\beta\;,\\
&\alpha AD+sB^2=B+\alpha\;, && \beta BC+tA^2=A+\beta\;,\\
&\beta AC+tAB=0\;, && \alpha BD+sAB=0\;,\\
&A+AC(A+B)=A\;, && B+BD(B+A)=B\;,\\
&A^2D+B^2C=A+B\;, && AB(C+D)=0\;,\\
&C+\alpha C^2+sAD=s\;, && D+\beta D^2+tBC=t\;,\\
&C(sA+\alpha D)=0\;, && D(tB+\beta C)=0\;,\\
&D(\alpha C+sB)=0\;, && C(\beta D+tA)=0\;,\\
&\beta C^2+tAD=t+C\;, && \alpha D^2+sBC=D+s\;,\\
&AC(C+D)=0\;, && BD(D+C)=0\;,\\
&DC(A+B)=0\;, && BC^2+AD^2=D+C\;.
\end{aligned}\tag{6.4}$$

While analyzing this system, we will assume that the blocks

$$(A,\alpha)\,,\qquad (B,\beta)\,,\qquad (D,t)\,,\qquad (C,s) \tag{6.5}$$

can be vanishing or not. The simplest possibility is

$$(A,\alpha)=0\,,\qquad (B,\beta)=0\,,\qquad (D,t)=0\,,\qquad (C,s)=0\,, \tag{6.6}$$

which leads to

solution $(KM-1)$,

$$G=\begin{pmatrix} K & 0 \\ 0 & M \end{pmatrix}\,. \tag{6.7}$$

One may separate 4 different cases with one non-vanishing block:

$$\begin{aligned}
&(A,\alpha)\neq 0\,,\quad (B,\beta)=0\,,\quad (D,t)=0\,,\quad (C,s)=0\,;\\
&(A,\alpha)=0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)=0\,,\quad (C,s)=0\,;\\
&(A,\alpha)=0\,,\quad (B,\beta)=0\,,\quad (D,t)\neq 0\,,\quad (C,s)=0\,;\\
&(A,\alpha)=0\,,\quad (B,\beta)=0\,,\quad (D,t)=0\,,\quad (C,s)\neq 0\,.
\end{aligned} \tag{6.8}$$

Moreover, there are six possible cases with two non-vanishing blocks (two of them marked by asterisk were considered in previous sections)

$$\begin{aligned}
&(A,\alpha)=0\,,\quad (B,\beta)=0\,,\quad (D,t)\neq 0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)=0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)=0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)=0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)\neq 0\,,\quad (C,s)=0\,;\qquad (*)\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)=0\,,\quad (D,t)\neq 0\,,\quad (C,s)=0\,;\qquad (*)\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)=0\,,\quad (D,t)=0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)=0\,,\quad (C,s)=0\,.
\end{aligned} \tag{6.9}$$

There exist four cases with three non-vanishing blocks:

$$\begin{aligned}
&(A,\alpha)=0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)\neq 0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)=0\,,\quad (D,t)\neq 0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)=0\,,\quad (C,s)\neq 0\,;\\
&(A,\alpha)\neq 0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)\neq 0\,,\quad (C,s)=0\,.
\end{aligned} \tag{6.10}$$

Finally, there is possible the case when all the four blocks are non-zero.
Let it be (see (6.8))

$$(A,\alpha)\neq 0\,,\quad (B,\beta)=0\,,\quad (D,t)=0\,,\quad (C,s)=0\,; \tag{6.11}$$

the system (6.4) becomes simpler and leads to the already known solution (6.7).
Let it be (see (6.8))

$$(A,\alpha)=0\,,\quad (B,\beta)\neq 0\,,\quad (D,t)=0\,,\quad (C,s)=0\,; \tag{6.12}$$

the system (6.4) again gives the known solution (6.7).
Let it be (see (6.8))

$$(A,\alpha)=0\,,\quad (B,\beta)=0\,,\quad (D,t)\neq 0\,,\quad (C,s)=0\,; \tag{6.13}$$

again we arrive at the solution (6.7).

Let it be (see (6.8))

$$(A,\alpha)=0\,,\qquad (B,\beta)=0\,,\qquad (D,t)=0\,,\qquad (C,s)\neq 0\,; \tag{6.14}$$

it leads to the known solution (6.7).

Now, let it be (see (6.9))

$$(A,\alpha)=0\,,\qquad (B,\beta)=0\,,\qquad (D,t)\neq 0\,,\qquad (C,s)\neq 0\,; \tag{6.15}$$

the system (6.4) takes the form

$$0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ 0=t+s\,,\ 0=0\,,\ 0=C+D\,,\ s=C\,,\ t=D\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ ;0=0\,,\ 0=0\,,\ ;C=s\,,\ D=t\,,\ ;0=0\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ 0=t+C\,,\ 0=D+s\,,\ 0=0\,,\ 0=0\,,\ ;0=0\,,\ 0=D+C\,,$$

with the following solution

$$t=D\,,\ s=C=-D\,,\ (A,\alpha)=0\,,\ (B,\beta)=0\,; \tag{6.16}$$

thus, we get

solution $(KM-2)$,

$$\mathbf{n}=0\,,\ n_0=0\,,\ \mathbf{l}=D(\mathbf{m}-\mathbf{k})\,,\ l_0=D(m_0-k_0)\,;$$

$$G=\begin{pmatrix} K & 0\\ D(M-K) & M\end{pmatrix};$$

$$G'G=\begin{pmatrix} K' & 0\\ D(M'-K') & M'\end{pmatrix}\begin{pmatrix} K'K & 0\\ D(M'M-K'K) & M'M\end{pmatrix}. \tag{6.17}$$

Now, let it be (see (6.9))

$$(A,\alpha)=0\,,\qquad (B,\beta)\neq 0\,,\qquad (D,t)=0\,,\qquad (C,s)\neq 0\,; \tag{6.18}$$

the system (6.4) reads

$$0=0\,,\ 0=0\,,\ \beta s=1\,,\ 0=0\,,\ 0=0\,,\ \beta C=1\,,\ 0=0\,,\ \beta=B\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ \beta s=1\,,\ 0=0\,,\ sB=1\,,\ s=C\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ B=\beta\,,\ sB=1\,,\ BC=1\,,\ ;0=0\,,\ 0=0\,,$$

$$0=0\,,\ B=B\,,\ BC=1\,,\ 0=0\,,\ C=s\,,0=0\,,\ ;0=0\,,\ 0=0\,,$$

$$0=0\,,\ \beta C=1\,,\ BC=1\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ BC=1\,,$$

which is equivalent to

$$s=C\,,\qquad \beta=B\,,\qquad BC=1\,. \tag{6.19}$$

Thus, we obtain

solution $(KM-3)$,

$$(A,\alpha)=0\,,\quad \beta=B\,,\quad (D,t)=0\,,\quad C=s=\frac{1}{B}\,;$$

$$\mathbf{n}=B\mathbf{m}\,,\quad n_0=Bm_0\,,\quad \mathbf{l}=\frac{1}{B}\mathbf{k}\,,\quad l_0=\frac{1}{B}k_0\,; \tag{6.20}$$

$$G=\begin{pmatrix} K & BM \\ B^{-1}K & M \end{pmatrix},\ G'G=\begin{pmatrix} (K'+M')K & B(K'+M')M \\ B^{-1}(K'+M')K & (K'+M')M \end{pmatrix}.$$

Let us consider the case (see (6.9))

$$(A,\alpha)\neq 0\,,\qquad (B,\beta)=0\,,\qquad (D,t)=0\,,\qquad (C,s)\neq 0\,; \tag{6.21}$$

the system (6.4) takes the form

$$\alpha s\alpha=0\,,\ 0=0\,,\ 0=\alpha\,,\ 0=0\,,\ 0=0\,,\ 0=A\,,\ \alpha+\alpha AC=A\,,\ 0=0\,,$$

$$s\alpha s=0\,,\ 0=0\,,\ 0=0\,,\ 0=s\,,\ 0=0\,,\ 0=C\,,\ s+ACs=C\,,\ 0=0\,,$$

$$A\alpha C=0\,,\ 0=0\,,\ A+sA^2=\alpha\,,\ 0=0\,,\ 0=\alpha\,,\ 0=A\,,\ 0=0\,,\ 0=0\,,$$

$$A+ACA=A\,,\ 0=0\,,\ 0=A\,,\ 0=0\,,\ C+\alpha C^2=s\,,\ 0=0\,,\ CsA=0\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ 0=C\,,\ 0=s\,,\ AC^2=0\,,\ 0=0\,,\ 0=0\,,\ 0=C\,,$$

which leads to already known

solution $(KM-1)$,

$$(B,\beta)=0\,,\ (D,t)=0\,,\ (A,\alpha)=0\,,\ (C,s)=0\,,\qquad G=\begin{pmatrix} K & 0 \\ 0 & M \end{pmatrix}. \tag{6.22}$$

Let us consider the case (see (6.9))

$$(A,\alpha)\neq 0\,,\qquad (B,\beta)\neq 0\,,\qquad (D,t)=0\,,\qquad (C,s)=0\,; \tag{6.23}$$

the system (6.4) gives

$$0=0\,,\ 0=0\,,\ 0=\beta+\alpha\,,\ 0=0\,,\ 0=0\,,\ 0=B+A\,,\ \alpha=A\,,\ \beta=B\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=D\,,$$

$$0=0\,,\ 0=0\,,\ A=\alpha\,,\ B=\beta\,,\ 0=B+\alpha\,,\ 0=A+\beta\,,\ 0=0\,,\ 0=0\,,$$

$$A=A\,,\ B=B\,,\ 0=A+B\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,$$

$$0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,\ 0=0\,,0=0\,,\ 0=0\,,\ 0=0\,,$$

which leads to

solution $(KM-4)$,

$$\begin{gathered} A=\alpha\,,\qquad B=\beta=-A\,,\qquad (D,t)=0\,,\qquad (C,s)=0\,,\\ \mathbf{n}=A(\mathbf{k}-\mathbf{m})\,,\ n_0=A(k_0-m_0)\,,\qquad \mathbf{l}=0\,,\ \ l_0=0\,, \end{gathered} \tag{6.24}$$

$$G=\begin{pmatrix} K & A(K-M) \\ 0 & M \end{pmatrix},\ G'G=\begin{pmatrix} K'K & A(K'K-M'M) \\ 0 & M'M \end{pmatrix}.$$

Now, we are able to consider the four possibilities with three vanishing blocks (see (6.10)). Let it be

$$(A,\alpha)=0\,,\qquad (B,\beta)\neq 0\,,\qquad (D,t)\neq 0\,,\qquad (C,s)\neq 0\,, \tag{6.25}$$

the system (6.4) gives

$$0=0\,,\ \beta t\beta=0\,,\ \beta^2 s=\beta\,,\ 0=0\,,\ 0=0\,,\ \beta CB=B\,,\ 0=0\,,\ \beta+\beta BD=B\,,$$

$$0=0\,,\ \beta t(s+t)=0\,,\ st\beta=0\,,\ \beta s^2=t+s\,,\ tCB=0\,,$$

$$sBC=C+D\,,\ s=C\,,\ t+BD(t+s)=D\,,$$

$$0=0\ ,\ B\beta D=0\ ,\ 0=0\ ,\ B+tB^2=\beta\ ,\ sB^2=B\ ,\ \beta BC=\beta\ ,\ 0=0\ ,\ 0=0\ ,$$
$$0=0\ ,\ B+BDB=B\ ,\ B^2C=B\ ,\ 0=0\ ,\ C=s\ ,\ D+\beta D^2+tBC=t\ ,$$
$$0=0\ ,\ (tB+\beta C)=0\ ,\ DsB=0\ ,\ C\beta D=0\ ,\ \beta C^2=t+C\ ,$$
$$sBC=D+s\ ,\ 0=0\ ,\ BD(D+C)=0\ ,\ DCB=0\ ,\ BC^2=D+C\ .$$

These equations lead to the already known solution $(KM-1)$.

For the case

$$(A,\alpha)\neq 0\ ,\qquad (B,\beta)=0\ ,\qquad (D,t)\neq 0\ ,\qquad (C,s)\neq 0 \tag{6.26}$$

the system (6.4) gives only one and already known solution $(KM-1)$.

Let it be

$$(A,\alpha)\neq 0\ ,\qquad (B,\beta)\neq 0\ ,\qquad (D,t)=0\ ,\qquad (C,s)\neq 0\ ; \tag{6.27}$$

the system (6.4) leads to the already known solution $(KM-1)$.

Consider the case where we assume

$$(A,\alpha)\neq 0\ ,\qquad (B,\beta)\neq 0\ ,\qquad (D,t)\neq 0\ ,\qquad (C,s)=0\ ; \tag{6.28}$$

the system (6.4) reads

$$0=0\ ,\ \beta t(\alpha+\beta)=0\ ,\ \alpha^2t=\beta+\alpha\ ,\ \alpha\beta t=0\ ,$$
$$\beta AD=0\ ,\ \alpha AD=B+A\ ,\ \alpha=A\ ,\ \beta+(\alpha+\beta)BD=B\ ,$$
$$0=0\ ,\ \beta tt=0\ ,\ 0=0\ ,\ \alpha t^2=t\ ,$$
$$0=0\ ,\ tAD=D\ ,\ 0=0\ ,\ t+BDt=D\ ,$$
$$0=0\ ,\ B(\beta D+tA)=0\ ,\ A=\alpha\ ,\ B+tB^2+\beta AD=\beta\ ,$$
$$\alpha AD=B+\alpha\ ,\ \beta tA^2=A+\beta\ ,\ tAB=0\ ,\ \alpha BD=0\ ,$$
$$A=A\ ,\ B+BD(B+A)=B\ ,\ A^2D=A+B\ ,\ ABD=0\ ,$$
$$0=0\ ,\ D+\beta D^2=t\ ,\ 0=0\ ,\ DtB=0\ ,$$
$$0=0\ ,\ 0=0\ ,\ tAD=t\ ,\ \alpha D^2=D\ ,$$
$$0=0\ ,\ BDD=0\ ,\ 0=0\ ,\ AD^2=D\ .$$

which leads to the already known solution $(KM-1)$.

It remains to examine the case of four non-zero blocks

$$(A,\alpha)\neq 0\ ,\qquad (B,\beta)\neq 0\ ,\qquad (D,t)\neq 0\ ,\qquad (C,s)\neq 0\ . \tag{6.29}$$

The system (6.4) can be simplified to

$$\begin{aligned}
&(\alpha+\beta)=0\ , && (\alpha+\beta)=0\ ,\\
&\alpha^2t+\beta^2s=\beta+\alpha\ , && (s+t)=0\ ,\\
&\alpha BC+\beta AD=0\ , && \alpha AD+\beta BC=B+A\ ,\\
&\alpha+(\alpha+\beta)AC=A\ , && \beta+(\alpha+\beta)BD=B\ ,
\end{aligned}$$

$$\begin{aligned}
&(s+t)=0\ , && (s+t)=0\ ,\\
&(\alpha+\beta)=0\ , && \beta s^2+\alpha t^2=t+s\ ,\\
&sAD+tCB=0, && sBC+tAD=C+D\ ,\\
&s+AC(s+t)=C\ , && t+BD(t+s)=D\ ,
\end{aligned}$$

$$
\begin{array}{ll}
(\alpha C + sB) = 0 \,, & (\beta D + tA) = 0 \,, \\
A + sA^2 + \alpha BC = \alpha \,, & B + tB^2 + \beta AD = \beta \,, \\
\alpha AD + sB^2 = B + \alpha \,, & \beta BC + tA^2 = A + \beta \,, \\
\beta C + tB = 0 \,, & \alpha D + sA = 0 \,, \\
(A + B) = 0 \,, & (B + A) = 0 \,, \\
A^2 D + B^2 C = A + B \,, & (C + D) = 0 \,, \\
C + \alpha C^2 + sAD = s \,, & D + \beta D^2 + tBC = t \,, \\
(sA + \alpha D) = 0 \,, & (tB + \beta C) = 0 \,,
\end{array}
$$

$$
\begin{array}{ll}
(\alpha C + sB) = 0 \,, & (\beta D + tA) = 0 \,, \\
\beta C^2 + tAD = t + C \,, & \alpha D^2 + sBC = D + s \,, \\
(C + D) = 0 \,, & (D + C) = 0 \,, \\
(A + B) = 0 \,, & BC^2 + AD^2 = D + C \,.
\end{array}
\tag{6.30}
$$

By eliminating the variables

$$
B = -A \,,\ \beta = -\alpha \,, \qquad C = -D \,,\ s = -t \,, \tag{6.31}
$$

we get the system

$$
\begin{array}{ll}
0 = 0 \,,\ 0 = 0 \,,\ 0 = 0 \,,\ 0 = 0 \,, & \\
0 = 0 \,,\ 0 = 0 \,,\ \alpha = A \,,\ \beta = -A \,, & \\
0 = 0 \,,\ 0 = 0 \,,\ 0 = 0 \,,\ 0 = 0 \,, & \\
0 = 0 \,,\ 0 = 0 \,,\ s = -D \,,\ t = D \,, & \\
(-\alpha D - sA) = 0 \,, & (\beta D + tA) = 0 \,, \\
A + sA^2 + \alpha AD = \alpha \,, & -A + tA^2 + \beta AD = \beta \,, \\
\alpha AD + sA^2 = -A + \alpha \,, & \beta AD + tA^2 = A + \beta \,, \\
\beta - D - tA = 0 \,, & \alpha D + sA = 0 \,, \\
0 = 0 \,, & 0 = 0 \,, \\
0 = 0 \,, & 0 = 0 \,, \\
-D + \alpha D^2 + sAD = s \,, & D + \beta D^2 + tAD = t \,, \\
(sA + \alpha D) = 0 \,, & (-tA - \beta D) = 0 \,, \\
(\alpha C - sA) = 0 \,, & (\beta D + tA) = 0 \,, \\
\beta C^2 + tAD = t + C \,, & \alpha D^2 - sAC = D + s \,, \\
0 = 0 \,, & 0 = 0 \,, \\
0 = 0 \,, & 0 = 0 \,.
\end{array}
$$

From this it follows

$$
A = \alpha \,, \qquad B = \beta = -A \,, \qquad t = D \,, \qquad C = s = -D \,, \tag{6.32}
$$

which describes the following

solution $(KM-5)$,

$$\begin{aligned}
\mathbf{n} &= A(\mathbf{k}-\mathbf{m})\,, \qquad n_0 = A(k_0-m_0)\,,\\
\mathbf{l} &= D(\mathbf{m}-\mathbf{k})\,, \qquad l_0 = D(m_0-k_0)\,;\\
G &= \begin{pmatrix} K & A(K-M) \\ C(K-M) & M \end{pmatrix}. \qquad (6.33)
\end{aligned}$$

Let us verify by direct calculation that the matrices with such structure form a group:

$$\begin{pmatrix} K' & A(K'-M') \\ C(K'-M') & M' \end{pmatrix}\begin{pmatrix} K & A(K-M) \\ C(K-M) & M \end{pmatrix} = \begin{pmatrix} K'' & A(K''-M'') \\ C(K''-M'') & M'' \end{pmatrix}.$$

7 Two Independent Vectors: Variant II(l,n)

Let us examine **variant II(l,n)**:

$$\begin{aligned}
\mathbf{k} &= (A\mathbf{l}+B\mathbf{n})\,, \qquad k_0 = (\alpha l_0+\beta n_0)\,,\\
\mathbf{m} &= (D\mathbf{l}+C\mathbf{n})\,, \qquad m_0 = (tl_0+sn_0)\,. \qquad (7.1)
\end{aligned}$$

The multiplication law takes the form

$$\begin{aligned}
k_0'' &= (\alpha l_0'+\beta n_0')\,(\alpha l_0+\beta n_0)+(A\mathbf{l}'+B\mathbf{n}')(A\mathbf{l}+B\mathbf{n})+n_0'\,l_0+\mathbf{n}'\cdot\mathbf{l}\,,\\
m_0'' &= (tl_0'+sn_0')\,(tl_0+sn_0)+(D\mathbf{l}'+C\mathbf{n}')\,(D\mathbf{l}+C\mathbf{n})+l_0'\,n_0+\mathbf{l}'\cdot\mathbf{n}\,,\\
n_0'' &= (\alpha l_0'+\beta n_0')\,n_0+(A\mathbf{l}'+B\mathbf{n}')\,\mathbf{n}+n_0'\,(tl_0+sn_0)+\mathbf{n}'\,(D\mathbf{l}+C\mathbf{n})\,,\\
l_0'' &= l_0'\,(\alpha l_0+\beta n_0)+\mathbf{l}'\,(A\mathbf{l}+B\mathbf{n})+(tl_0'+sn_0')\,l_0+(D\mathbf{l}'+C\mathbf{n}')\,\mathbf{l}\,,
\end{aligned}$$

$$\begin{aligned}
\mathbf{k}'' &= (\alpha l_0'+\beta n_0')\,(A\mathbf{l}+B\mathbf{n})+(A\mathbf{l}'+B\mathbf{n}')\,(\alpha l_0+\beta n_0)\\
&+i\,(A\mathbf{l}'+B\mathbf{n}')\times(A\mathbf{l}+B\mathbf{n})+n_0'\,\mathbf{l}+\mathbf{n}'\,l_0+i\,\mathbf{n}'\times\mathbf{l}\,,\\
\mathbf{m}'' &= (tl_0'+sn_0')\,(D\mathbf{l}+C\mathbf{n})+(D\mathbf{l}'+C\mathbf{n}')\,(tl_0+sn_0)\\
&+i\,(D\mathbf{l}'+C\mathbf{n}')\times(D\mathbf{l}+C\mathbf{n})+l_0'\,\mathbf{n}+\mathbf{l}'\,n_0+i\,\mathbf{l}'\times\mathbf{n}\,,\\
\mathbf{n}'' &= (\alpha l_0'+\beta n_0')\,\mathbf{n}+(A\mathbf{l}'+B\mathbf{n}')\,n_0+i\,(A\mathbf{l}'+B\mathbf{n}')\times\mathbf{n}\\
&+n_0'\,(D\mathbf{l}+C\mathbf{n})+\mathbf{n}'\,(tl_0+sn_0)+i\,\mathbf{n}'\times(D\mathbf{l}+C\mathbf{n})\,,
\end{aligned}$$

$$\begin{aligned}
\mathbf{l}'' &= l_0'\,(A\mathbf{l}+B\mathbf{n})+\mathbf{l}'\,(\alpha l_0+\beta n_0)+i\,\mathbf{l}'\times(A\mathbf{l}+B\mathbf{n})\\
&+(tl_0'+sn_0')\,\mathbf{l}+(D\mathbf{l}'+C\mathbf{n}')\,l_0+i\,(D\mathbf{l}'+C\mathbf{n}')\times\mathbf{l}\,. \qquad (7.2)
\end{aligned}$$

The equation

$$\mathbf{k}'' = A\mathbf{l}''+B\mathbf{n}'' \qquad (7.3)$$

takes the form

$$\begin{aligned}
&(\alpha l_0'+\beta n_0')(A\mathbf{l}+B\mathbf{n})+(A\mathbf{l}'+B\mathbf{n}')(\alpha l_0+\beta n_0)\\
&+i(A\mathbf{l}'+B\mathbf{n}')\times(A\mathbf{l}+B\mathbf{n})+n_0'\mathbf{l}+\mathbf{n}'l_0+i\mathbf{n}'\times\mathbf{l}\\
&=A\,[l_0'(A\mathbf{l}+B\mathbf{n})+\mathbf{l}'(\alpha l_0+\beta n_0)+i\mathbf{l}'\times(A\mathbf{l}+B\mathbf{n})\\
&+(tl_0'+sn_0')\mathbf{l}+(D\mathbf{l}'+C\mathbf{n}')l_0+i(D\mathbf{l}'+C\mathbf{n}')\times\mathbf{l}]\\
&+B\,[(\alpha l_0'+\beta n_0')\mathbf{n}+(A\mathbf{l}'+B\mathbf{n}')n_0+i(A\mathbf{l}'+B\mathbf{n}')\times\mathbf{n}\\
&+n_0'(D\mathbf{l}+C\mathbf{n})+\mathbf{n}'\,(tl_0+sn_0)+i\mathbf{n}'\times(D\mathbf{l}+C\mathbf{n})]\,;
\end{aligned}$$

whence it follows

$$\begin{aligned}
l_0'\mathbf{l}\mathbf{l}: \quad & \alpha A = A^2 + At\,, \quad n_0'\mathbf{n}: \quad \beta B = \beta B + BC\,,\\
n_0'\mathbf{l}: \quad & \beta A + 1 = As + BD\,, \quad l_0'\mathbf{n}: \quad \alpha B = AB + \alpha B\,,\\
n_0\mathbf{l}': \quad & \beta A = \beta A + BA\,, \quad l_0\mathbf{n}': \quad \alpha B + 1 = AC + Bt\,,\\
l_0\mathbf{l}': \quad & \alpha A = \alpha A + AD\,, \quad n_0\mathbf{n}': \quad \beta B = B^2 + Bs\,,\\
\mathbf{l}'\times\mathbf{l}: \quad & A^2 = A^2 + AD\,, \quad \mathbf{n}'\times\mathbf{n}: \quad B^2 = B^2 + BC\,,\\
\mathbf{n}'\times\mathbf{l}: \quad & BA + 1 = AC + BD\,, \quad \mathbf{l}'\times\mathbf{n}: \quad AB = AB + AB\,.
\end{aligned} \tag{7.4}$$

The equation

$$\mathbf{m}'' = D\mathbf{l}'' + C\mathbf{n}'' \tag{7.5}$$

takes the form

$$\begin{aligned}
&(tl_0' + sn_0')(D\mathbf{l} + C\mathbf{n}) + (D\mathbf{l}' + C\mathbf{n}')(tl_0 + sn_0)\\
&+i(D\mathbf{l}' + C\mathbf{n}')\times(D\mathbf{l} + C\mathbf{n}) + l_0'\mathbf{n} + \mathbf{l}'n_0 + i\,\mathbf{l}'\cdot\mathbf{n} =\\
&\times D\,[l_0'(A\mathbf{l} + B\mathbf{n}) + \mathbf{l}'(\alpha l_0 + \beta n_0) + i\mathbf{l}'\times(A\mathbf{l} + B\mathbf{n})\\
&+(tl_0' + sn_0')\mathbf{l} + (D\mathbf{l}' + C\mathbf{n}')l_0 + i(D\mathbf{l}' + C\mathbf{n}')\times\mathbf{l}]\\
&+C\,[\alpha l_0' + \beta n_0')\,\mathbf{n} + (A\mathbf{l}' + B\mathbf{n}')\,n_0 + i\,(A\mathbf{l}' + B\mathbf{n}')\times\mathbf{n}\\
&+n_0'\,(D\mathbf{l} + C\mathbf{n}) + \mathbf{n}'\,(tl_0 + sn_0) + i\,\mathbf{n}'\times(D\mathbf{l} + C\mathbf{n})\,,
\end{aligned}$$

from which we obtain

$$\begin{aligned}
l_0'\mathbf{l}: \quad & tD = DA + tD\,, \quad n_0'\mathbf{n}: \quad sC = C\beta + C^2\,,\\
n_0'\mathbf{l}: \quad & sD = sD + CD\,, \quad l_0'\mathbf{n}: \quad tC + 1 = DB + \alpha C\,,\\
n_0\mathbf{l}': \quad & sD + 1 = \beta D + CA\,, \quad l_0\mathbf{n}': \quad tC = DC + tC\,,\\
l_0\mathbf{l}': \quad & tD = \alpha D + D^2\,, \quad n_0\mathbf{n}': \quad sC = CB + sC\,,\\
\mathbf{l}'\times\mathbf{l}: \quad & D^2 = DA + D^2\,, \quad \mathbf{n}'\times\mathbf{n}: \quad C^2 = CB + C^2\,,\\
\mathbf{n}'\times\mathbf{l}: \quad & CD = CD + CD\,, \quad \mathbf{l}'\times\mathbf{n}: \quad CD + 1 = DB + CA\,.
\end{aligned}$$

The equation

$$k_0'' = \alpha l_0'' + \beta n_0'' \tag{7.6}$$

gives

$$\begin{aligned}
&(\alpha l_0' + \beta n_0')\,(\alpha l_0 + \beta n_0) + (A\mathbf{l}' + B\mathbf{n}')(A\mathbf{l} + B\mathbf{n}) + n_0'\,l_0 + \mathbf{n}'\cdot\mathbf{l}\\
&= \alpha\,[\,l_0'\,(\alpha l_0 + \beta n_0) + \mathbf{l}'\,(A\mathbf{l} + B\mathbf{n}) + (tl_0' + sn_0')\,l_0 + (D\mathbf{l}' + C\mathbf{n}')\,\mathbf{l}]\\
&+\beta\,[\,(\alpha l_0' + \beta n_0')\,n_0 + (A\mathbf{l}' + B\mathbf{n}')\,\mathbf{n} + n_0'\,(tl_0 + sn_0) + \mathbf{n}'\,(D\mathbf{l} + C\mathbf{n})]\,,
\end{aligned}$$

which infers

$$\begin{aligned}
l_0'l_0: \quad & \alpha^2 = \alpha^2 + \alpha t\,, \quad n_0'n_0: \quad \beta^2 = \beta^2 + s\beta\,,\\
l_0'n_0: \quad & \alpha\beta = \alpha\beta + \alpha\beta\,, \quad n_0'l_0: \quad \alpha\beta + 1 = \alpha s + \beta t\,,\\
\mathbf{l}'\cdot\mathbf{l}: \quad & A^2 = \alpha A + \alpha D\,, \quad \mathbf{n}'\cdot\mathbf{n}: \quad B^2 = \beta B + \beta C\,,\\
\mathbf{l}'\cdot\mathbf{n}: \quad & AB = \alpha B + \beta A\,, \quad \mathbf{n}'\cdot\mathbf{l}: \quad AB + 1 = \alpha C + \beta D\,.
\end{aligned}$$

The equation

$$m_0'' = tl_0'' + sn_0'' \tag{7.7}$$

takes the form

$$\begin{aligned}
&(tl_0' + sn_0')\,(tl_0 + sn_0) + (D\mathbf{l}' + C\mathbf{n}')\,(D\mathbf{l} + C\mathbf{n}) + l_0'\,n_0 + \mathbf{l}' \cdot \mathbf{n} \\
&= t\,[\; l_0'\,(\alpha l_0 + \beta n_0) + \mathbf{l}'\,(A\mathbf{l} + B\mathbf{n}) + (tl_0' + sn_0')\,l_0 + (D\mathbf{l}' + C\mathbf{n}')\,\mathbf{l}\;] \\
&+ s\,[\; (\alpha l_0' + \beta n_0')\,n_0 + (A\mathbf{l}' + B\mathbf{n}')\,\mathbf{n} + n_0'\,(tl_0 + sn_0) + \mathbf{n}'\,(D\mathbf{l} + C\mathbf{n})\;]\;,
\end{aligned}$$

from which we obtain

$$\begin{aligned}
l_0'l_0: &\qquad t^2 = t\alpha + t^2\;, \quad n_0'n_0: \qquad s^2 = s\beta + s^2\;, \\
l_0'n_0: &\qquad ts + 1 = t\beta + s\alpha\;, \quad n_0'l_0: \qquad st = st + st\;, \\
\mathbf{l}'\cdot\mathbf{l}: &\qquad D^2 = tA + tD\;, \quad \mathbf{n}'\cdot\mathbf{n}: \qquad C^2 = sB + sC\;, \\
\mathbf{l}'\cdot\mathbf{n}: &\qquad DC + 1 = tB + sA\;, \quad \mathbf{n}'\cdot\mathbf{l}: \qquad CD = tC + sD\;.
\end{aligned}$$

We further collect the equations together, and infer

$$\alpha A = A^2 + At\;,\; 0 = BC\;,\; \beta A + 1 = As + BD\;,\; 0 = AB\;,\; 0 = +BA\;,$$

$$\alpha B + 1 = AC + Bt\;,\; 0 = AD\;,\; \beta B = B^2 + Bs\;,\; 0 = AD\;,\; 0 = BC\;,$$

$$BA + 1 = AC + BD\;,\; 0 = AB\;,\; 0 = DA\;,\; sC = C\beta + C^2\;,\; 0 = CD\;,$$

$$tC + 1 = DB + \alpha C\;,\; sD + 1 = \beta D + CA\;,\; 0 = DC\;,\; tD = \alpha D + D^2\;,\; 0 = CB\;,$$

$$0 = DA\;,\; 0 = CB\;,\; 0 = CD\;,\; CD + 1 = DB + CA\;;$$

$$0 = \alpha t\;, \quad 0 = s\beta\;, \quad 0 = \alpha\beta\;, \quad \alpha\beta + 1 = \alpha s + \beta t\;,$$

$$A^2 = \alpha A + \alpha D\;,\; B^2 = \beta B + \beta C\;,\; AB = \alpha B + \beta A\;,\; AB + 1 = \alpha C + \beta D\;;$$

$$0 = t\alpha\;, \quad 0 = s\beta\;, \quad ts + 1 = t\beta + s\alpha\;, \quad 0 = st\;,$$

$$D^2 = tA + tD\;,\; C^2 = sB + sC\;,\; DC + 1 = tB + sA\;,\; CD = tC + sD\;.$$

From the very beginning we note that the trivial set of all vanishing parameters provide no solution of the system.

Let us separate the most simple equations

$$\begin{aligned}
&BC = 0\;, \quad \beta s = 0\;, \quad AB = 0\;, \quad \alpha\beta = 0\;, \\
&AD = 0\;, \quad \alpha t = 0\;, \quad CD = 0\;, \quad st = 0\;;
\end{aligned} \tag{7.8}$$

they can be satisfied by six solutions

$$\begin{aligned}
&1) \quad B, \beta = 0\;, \quad D, t = 0\;; \quad 2) \quad A, \alpha = 0\;, \quad C, s = 0\;; \\
&3) \qquad B, \beta = 0\;, \qquad A, \alpha = 0\;, \qquad C, s = 0\;; \\
&4) \qquad A, \alpha = 0\;, \qquad B, \beta = 0\;, \qquad D, t = 0\;; \\
&5) \qquad C, s = 0\;, \qquad B, \beta = 0\;, \qquad D, t = 0\;; \\
&6) \qquad D, t = 0\;, \qquad C, s = 0\;, \qquad A, \alpha = 0\;.
\end{aligned} \tag{7.9}$$

The remaining equations are

$$\alpha A = A^2 + At\ , \quad \beta A + 1 = As + BD\ , \quad \alpha B + 1 = AC + Bt\ ,$$
$$\beta B = B^2 + Bs\ , \quad BA + 1 = AC + BD\ ; \tag{7.10}$$

$$sC = C\beta + C^2\ , \quad tC + 1 = DB + \alpha C\ , \quad sD + 1 = \beta D + CA\ ,$$
$$tD = \alpha D + D^2\ , \quad CD + 1 = DB + CA\ ; \tag{7.11}$$

$$\alpha\beta + 1 = \alpha s + \beta t\ , \quad A^2 = \alpha A + \alpha D\ , \quad B^2 = \beta B + \beta C\ ,$$
$$AB = \alpha B + \beta A\ , \quad AB + 1 = \alpha C + \beta D\ ; \tag{7.12}$$

$$ts + 1 = t\beta + s\alpha\ , \quad D^2 = tA + tD\ , \quad C^2 = sB + sC\ ,$$
$$DC + 1 = tB + sA\ , \quad CD = tC + sD\ . \tag{7.13}$$

We see that variants 3), 4), 5), 6) are not consistent with the additional equations (7.10)–(7.13). Therefore, we shall further examine only the variants 1) and 2).

In the case 1), we have

$$1) \qquad B, \beta = 0\ , \qquad D, t = 0\ ;$$

the equations (7.10)–(7.13) give

$$\begin{array}{ccc} \alpha A = A^2\ , & 1 = As\ , & 1 = AC\ , \\ sC = C^2\ , & 1 = \alpha C\ , & 1 = CA\ , \\ 1 = \alpha s\ , & A^2 = \alpha A\ , & 1 = \alpha C\ , \\ 1 = s\alpha\ , & C^2 = sC\ , & 1 = sA\ ; \end{array}$$

the last system has only one solution

$$B, \beta = 0\ , \qquad D, t = 0\ , \qquad \alpha = A \neq 0\ , \qquad C = s = \frac{1}{A}\ ; \tag{7.14}$$

so we arrive at

solution $(LN-1)$,

$$\mathbf{k} = A\mathbf{l}\ ,\ k_0 = Al_0\ ,\ \mathbf{m} = A^{-1}\mathbf{n}\ ,\ m_0 = A^{-1}n_0\ ,$$
$$G = \begin{pmatrix} AL & N \\ L & A^{-1}N \end{pmatrix}, \tag{7.15}$$
$$G'G = \begin{pmatrix} A(AL'L + A^{-1}N'L) & (AL'N + A^{-1}N'N) \\ (AL'L + A^{-1}N'L) & A^{-1}(AL'N + A^{-1}N'N) \end{pmatrix}.$$

This is a set of degenerate matrices of rank 2.

For the case

$$2) \qquad A, \alpha = 0\ , \qquad C, s = 0$$

the equations (7.10)–(7.13) give

$$\begin{array}{ccc} 1 = BD\ , & 1 = Bt\ , & \beta B = B^2\ , \\ 1 = DB\ , & 1 = \beta D\ , & tD = D^2\ , \\ 1 = \beta t\ , & B^2 = \beta B\ , & 1 = \beta D\ , \\ 1 = t\beta\ , & D^2 = tD\ , & 1 = tB\ , \end{array}$$

which has only one solution

$$A, \alpha = 0\ , \qquad C, s = 0\ , \qquad \beta = \beta \neq 0\ , \qquad D = t = B^{-1}\ ; \tag{7.16}$$

so we obtain

solution $(LN-2)$,

$$\begin{gathered}
\mathbf{k} = B\mathbf{n}\ ,\ k_0 = Bn_0\ ,\ \mathbf{m} = B^{-1}\mathbf{l}\ ,\ m_0 = B^{-1}l_0\ , \\
G = \begin{pmatrix} BN & N \\ L & B^{-1}L \end{pmatrix}, \\
G'G = \begin{pmatrix} B(BN'N + B^{-1}N'L) & (BN'N + B^{-1}N'L) \\ (BL'N + B^{-1}L'L) & B^{-1}(BL'N + B^{-1}L'L) \end{pmatrix}.
\end{gathered} \tag{7.17}$$

This is a set of degenerate matrices of rank 2. The analysis of the **variant II(l,n)** is completed.

8 Two Independent Vectors: Variant II(k,l)

Let us examine the **variant II(k,l)**:

$$\begin{aligned}
\mathbf{n} &= A\mathbf{k} + B\mathbf{l}\ , \qquad & n_0 &= \alpha k_0 + \beta l_0\ , \\
\mathbf{m} &= D\mathbf{l} + C\mathbf{k}\ , \qquad & m_0 &= tl_0 + sk_0\ .
\end{aligned} \tag{8.1}$$

The multiplication law takes the form

$$\begin{aligned}
k_0'' &= k_0'\, k_0 + \mathbf{k}' \cdot \mathbf{k} + (\alpha k_0' + \beta l_0')\, l_0 + (A\mathbf{k}' + B\mathbf{l}')\, \mathbf{l}\ , \\
m_0'' &= (tl_0' + sk_0')(tl_0 + sk_0) + (D\mathbf{l}' + C\mathbf{k}')\ (D\mathbf{l} + C\mathbf{k}) \\
&\quad + l_0'\ (\alpha k_0 + \beta l_0) + \mathbf{l}'(A\mathbf{k} + B\mathbf{l})\ , \\
n_0'' &= k_0'\ (\alpha k_0 + \beta l_0) + \mathbf{k}'\ (A\mathbf{k} + B\mathbf{l}) \\
&\quad + (\alpha k_0' + \beta l_0')\ (tl_0 + sk_0) + (A\mathbf{k}' + B\mathbf{l}')\ (D\mathbf{l} + C\mathbf{k})\ , \\
l_0'' &= l_0'\, k_0 + \mathbf{l}' \cdot \mathbf{k} + (tl_0' + sk_0')l_0 + (D\mathbf{l}' + C\mathbf{k}')\, \mathbf{l}\ ,
\end{aligned}$$

$$\begin{aligned}
\mathbf{k}'' &= k_0'\, \mathbf{k} + \mathbf{k}'\, k_0 + i\, \mathbf{k}' \times \mathbf{k} + (\alpha k_0' + \beta l_0')\, \mathbf{l} + (A\mathbf{k}' \\
&\quad + B\mathbf{l}')\, l_0 + i\ (A\mathbf{k}' + B\mathbf{l}') \times \mathbf{l}\ , \\
\mathbf{m}'' &= (tl_0' + sk_0')(D\mathbf{l} + C\mathbf{k}) + (D\mathbf{l}' + C\mathbf{k}')\ (tl_0 + sk_0) \\
&\quad + i\ (D\mathbf{l}' + C\mathbf{k}') \times (D\mathbf{l} + C\mathbf{k}) \\
&\quad + l_0'\ (A\mathbf{k} + B\mathbf{l}) + \mathbf{l}'\ (\alpha k_0 + \beta l_0) + i\, \mathbf{l}' \times (A\mathbf{k} + B\mathbf{l})\ , \\
\mathbf{n}'' &= k_0'\ (A\mathbf{k} + B\mathbf{l}) + \mathbf{k}'\ (\alpha k_0 + \beta l_0) + i\, \mathbf{k}' \times (A\mathbf{k} + B\mathbf{l}) \\
&\quad + (\alpha k_0' + \beta l_0')\ (D\mathbf{l} + C\mathbf{k}) \\
&\quad + (A\mathbf{k}' + B\mathbf{l}')\ (tl_0 + sk_0) + i\ (A\mathbf{k}' + B\mathbf{l}') \times (D\mathbf{l} + C\mathbf{k})\ , \\
\mathbf{l}'' &= l_0'\, \mathbf{k} + \mathbf{l}'\, k_0 + i\, \mathbf{l}' \times \mathbf{k} \\
&\quad + (tl_0' + sk_0')\mathbf{l} + (D\mathbf{l}' + C\mathbf{k}')\, l_0 + i\ (D\mathbf{l}' + C\mathbf{k}') \times \mathbf{l}\ .
\end{aligned} \tag{8.2}$$

The equation

$$\mathbf{n}'' = A\mathbf{k}'' + B\mathbf{l}'' \tag{8.3}$$

gives

$$\begin{aligned}
&k_0'\ (A\mathbf{k} + B\mathbf{l}) + \mathbf{k}'\ (\alpha k_0 + \beta l_0) + i\ \mathbf{k}' \times (A\mathbf{k} + B\mathbf{l}) \\
&+ (\alpha k_0' + \beta l_0')\ (D\mathbf{l} + C\mathbf{k}) + (A\mathbf{k}' + B\mathbf{l}')\ (tl_0 + sk_0) + i\ (A\mathbf{k}' + B\mathbf{l}') \times (D\mathbf{l} + C\mathbf{k}) \\
&= A\ [\ k_0'\, \mathbf{k} + \mathbf{k}'\, k_0 + i\, \mathbf{k}' \times \mathbf{k} + (\alpha k_0' + \beta l_0')\, \mathbf{l} + (A\mathbf{k}' + B\mathbf{l}')\, l_0 + i\ (A\mathbf{k}' + B\mathbf{l}') \times \mathbf{l}\ ] \\
&\quad + B\ [\ l_0'\, \mathbf{k} + \mathbf{l}'\, k_0 + i\, \mathbf{l}' \times \mathbf{k} + (tl_0' + sk_0')\mathbf{l} + (D\mathbf{l}' + C\mathbf{k}')\, l_0 + i\ (D\mathbf{l}' + C\mathbf{k}') \times \mathbf{l}\ ].
\end{aligned} \tag{8.4}$$

Further, we obtain the system

$$k_0' \, \mathbf{k} : \qquad \alpha C = 0 \; , \quad k_0 \, \mathbf{k}' : \qquad \alpha + As = A \; ,$$

$$k_0' \, \mathbf{l} : \qquad B + \alpha D = A\alpha + Bs \; , \quad k_0 \, \mathbf{l}' : \qquad sB = B \; ,$$

$$l_0' \, \mathbf{l} : \qquad \beta D = A\beta + Bt \; , \quad l_0 \, \mathbf{l}' : \qquad Bt = AB + BD \; ,$$

$$l_0' \, \mathbf{k} : \qquad \beta C = B \; , \quad l_0 \, \mathbf{k}' : \qquad \beta + At = A^2 + BC \; ,$$

$$\mathbf{k}' \times \mathbf{k} : \qquad AC = 0 \; , \quad \mathbf{l}' \times \mathbf{l} : \qquad AB = 0 \; ,$$

$$\mathbf{k}' \times \mathbf{l} : \qquad B + AD = A^2 + BC \; , \quad \mathbf{l}' \times \mathbf{k} : \qquad BC = B \; .$$

The equation

$$n_0'' = \alpha k_0'' + \beta l_0''$$

leads to

$$\begin{aligned} & k_0' \, (\alpha k_0 + \beta l_0) + \mathbf{k}' \, (A\mathbf{k} + B\mathbf{l}) \\ & + (\alpha k_0' + \beta l_0') \, (t l_0 + s k_0) + (A\mathbf{k}' + B\mathbf{l}') \, (D\mathbf{l} + C\mathbf{k}) = \\ & \alpha \, [\, k_0' \, k_0 + \mathbf{k}' \, \mathbf{k} + (\alpha k_0' + \beta l_0') \, l_0 + (A\mathbf{k}' + B\mathbf{l}') \, \mathbf{l} \,] \\ & + \beta \, [\, l_0' \, k_0 + \mathbf{l}' \cdot \mathbf{k} + (t l_0' + s k_0') l_0 + (D\mathbf{l}' + C\mathbf{k}') \cdot \mathbf{l} \,] \; ; \end{aligned}$$

this infers

$$k_0' k_0 : \qquad \alpha s = 0 \; , \quad l_0' l_0 : \qquad 0 = \alpha\beta \; ,$$

$$k_0' l_0 : \qquad \beta + \alpha t = \alpha^2 + \beta s \; , \quad l_0' k_0 : \qquad \beta s = \beta \; ,$$

$$\mathbf{k}' \cdot \mathbf{k} : \qquad A + AC = \alpha \; , \quad \mathbf{l}' \cdot \mathbf{l} : \qquad BD = \alpha B + \beta D \; ,$$

$$\mathbf{k}' \cdot \mathbf{l} : \qquad B + AD = \alpha A + \beta C \; , \quad \mathbf{k} \cdot \mathbf{l}' : \qquad BC = \beta \; .$$

The equation

$$\mathbf{m}'' = D\mathbf{l}'' + C\mathbf{k}'' \tag{8.5}$$

leads to

$$\begin{aligned} & (t l_0' + s k_0')(D\mathbf{l} + C\mathbf{k}) + (D\mathbf{l}' + C\mathbf{k}') \, (t l_0 + s k_0) + i \, (D\mathbf{l}' + C\mathbf{k}') \times (D\mathbf{l} + C\mathbf{k}) \\ & + l_0' \, (A\mathbf{k} + B\mathbf{l}) + \mathbf{l}' \, (\alpha k_0 + \beta l_0) + i \, \mathbf{l}' \times (A\mathbf{k} + B\mathbf{l}) = D \, [\, l_0' \, \mathbf{k} + \mathbf{l}' \, k_0 + i \, \mathbf{l}' \times \mathbf{k} \\ & + (t l_0' + s k_0')\mathbf{l} + (D\mathbf{l}' + C\mathbf{k}') \, l_0 + i \, (D\mathbf{l}' + C\mathbf{k}') \times \mathbf{l} \,] + C \, [\, k_0' \, \mathbf{k} + \mathbf{k}' \, k_0 + i \, \mathbf{k}' \times \mathbf{k} \\ & + (\alpha k_0' + \beta l_0') \, \mathbf{l} + (A\mathbf{k}' + B\mathbf{l}') \, l_0 + i \, (A\mathbf{k}' + B\mathbf{l}') \times \mathbf{l} \,] \; ; \end{aligned}$$

further we obtain

$$k_0' \, \mathbf{k} : \qquad sC = C \; , \quad k_0 \, \mathbf{k}' : \qquad sC = C \; ,$$

$$k_0' \, \mathbf{l} : \qquad sD = sD + C\alpha \; , \quad k_0 \, \mathbf{l}' : \qquad Ds + \alpha = D \; ,$$

$$l_0' \, \mathbf{l} : \qquad tD + B = Dt + C\beta \; , \quad l_0 \, \mathbf{l}' : \qquad Dt + \beta = D^2 + CB \; ,$$

$$0 \, \mathbf{k} : \qquad tC + A = D \; , \quad l_0 \, \mathbf{k}' : \qquad tC = DC + CA \; ,$$

$$\mathbf{k}' \times \mathbf{k} : \qquad C^2 = C \; , \quad \mathbf{l}' \times \mathbf{l} : \qquad D^2 + B = D^2 + CB \; ,$$

$$\mathbf{k}' \times \mathbf{l} : \qquad CD = CD + CA \; , \quad \mathbf{l}' \times \mathbf{k} : \qquad DC + A = D \; .$$

The equation

$$m_0'' = t l_0'' + s k_0'' \tag{8.6}$$

gives

$$(tl'_0 + sk'_0)(tl_0 + sk_0) + (D\mathbf{l}' + C\mathbf{k}')\,(D\mathbf{l} + C\mathbf{k}) + l'_0\,(\alpha k_0 + \beta l_0) + \mathbf{l}'(A\mathbf{k} + B\mathbf{l})$$
$$= t\,[\,l'_0\,k_0 + \mathbf{l}'\,\mathbf{k} + (tl'_0 + sk'_0)l_0 + (D\mathbf{l}' + C\mathbf{k}')\,\mathbf{l}\,]$$
$$+s\,[\,k'_0\,k_0 + \mathbf{k}'\cdot\mathbf{k} + (\alpha k'_0 + \beta l'_0)\,l_0 + (A\mathbf{k}' + B\mathbf{l}')\cdot\mathbf{l}\,]\,;$$

whence it follows

$$k'_0k_0: \quad s^2 = s\,, \quad l'_0l_0: \quad t^2 + \beta = t^2 + s\beta\,,$$
$$k'_0l_0: \quad st = ts + s\alpha\,, \quad l'_0k_0: \quad ts + \alpha = t\,,$$
$$\mathbf{k}'\cdot\mathbf{k}: \quad C^2 = s\,, \quad \mathbf{l}'\cdot\mathbf{l}: \quad D^2 + B = tD + sB\,,$$
$$\mathbf{k}'\cdot\mathbf{l}: \quad CD = tC + sA\,, \quad \mathbf{k}\cdot\mathbf{l}': \quad DC + A = t\,.$$

Let us collect the equations together:

$$\alpha C = 0\,, \quad AC = 0\,, \quad AB = 0\,,$$
$$(s-1)B = 0\,, \quad \beta C = B\,, \quad (C-1)B = 0\,,$$
$$\alpha + A(s-1) = 0\,, \quad (A-D)\alpha + B(s-1) = 0\,,$$
$$(A-D)\beta + Bt = 0\,, \quad AB + B(D-t) = 0\,,$$
$$\beta + At = A^2 + BC\,, \quad B + AD = A^2 + BC\,, \tag{8.7}$$

$$\alpha s = 0\,, \quad \alpha\beta = 0\,,$$
$$(s-1)\beta = 0\,, \quad BC = \beta\,,$$
$$\alpha t = \alpha^2 + \beta(s-1)\,, \quad A + AC = \alpha\,,$$
$$(B-\beta)D = \alpha B\,, \quad B + AD = \alpha A + \beta C\,, \tag{8.8}$$

$$C\alpha = 0\,, \quad CA = 0\,,$$
$$(s-1)C = 0\,, \quad B = C\beta\,, \quad C(C-1) = 0\,, \quad (C-1)B = 0\,,$$
$$D(s-1) + \alpha = 0\,, \quad Dt + \beta = D^2 + CB\,,$$
$$tC + A = D\,, \quad (D-t)C + CA = 0\,, \quad DC + A = D\,; \tag{8.9}$$

$$0 = s\alpha\,, \quad s(s-1) = 0\,, \quad (s-1)\beta = 0\,,$$
$$t(s-1) + \alpha = 0\,, \quad C^2 = s\,,$$
$$D(D-t) = (s-1)B\,, \quad C(D-C) = sA\,, \quad DC + A = t\,. \tag{8.10}$$

For s, only two values are possible: $s = 0$, 1. For $s = 0$, the system has only one solution, namely

solution $(KL-1)$,

$$\alpha = A\,, \quad B = \beta = 0\,, \quad C = s = 0\,, \quad D = t = A\,,$$
$$\mathbf{n} = A\mathbf{k}\,, \quad n_0 = Ak_0\,, \quad \mathbf{m} = A\mathbf{l}\,, \quad m_0 = Al_0\,, \tag{8.11}$$
$$G = \begin{pmatrix} K & AK \\ L & AL \end{pmatrix}, \quad G'G = \begin{pmatrix} (K'K + AK'L) & A(K'K + AK'L) \\ (L'K + AL'L) & A(L'K + AL'L) \end{pmatrix}.$$

For $s = 1$, the system also has only one solution:

solution $(KL-2)$,

$$C = s = +1\,, \quad A = \alpha = 0\,, \quad D = t = +1\,, \quad B = \beta = 1\,,$$
$$\mathbf{n} = \mathbf{l}\,, \quad n_0 = l_0\,, \quad \mathbf{m} = \mathbf{l} + \mathbf{k}\,, \quad m_0 = l_0 + k_0\,, \tag{8.12}$$
$$G = \begin{pmatrix} K & L \\ L & K + L \end{pmatrix},$$
$$G'G = \begin{pmatrix} (K'K + L'L) & (L'K + K'L + L'L) \\ (L'K + K'L + L'L) & [(K'K + L'L) + (L'K + K'L + L'L)] \end{pmatrix}.$$

9 Two Independent Vectors: Variant I(n,m)

Let us examine **variant II(n,m)**:

$$\begin{aligned} \mathbf{l} = A\mathbf{m} + B\mathbf{n}\ , \qquad & n_0 = \alpha m_0 + \beta n_0\ , \\ \mathbf{k} = D\mathbf{n} + C\mathbf{m}\ , \qquad & k_0 = t n_0 + s m_0\ . \end{aligned} \tag{9.1}$$

Here we get only

solution $(NM-1)$,

$$G = \begin{pmatrix} AN & N \\ AM & M \end{pmatrix}; \tag{9.2}$$

solution $(NM-2)$,

$$G = \begin{pmatrix} M+N & N \\ N & M \end{pmatrix}. \tag{9.3}$$

10 Two Independent Vectors: Variant II(k,n)

Let us examine **variant II(k,n)**:

$$\begin{aligned} \mathbf{l} = (A\mathbf{k} + B\mathbf{n})\ , \qquad & l_0 = (\alpha k_0 + \beta n_0)\ , \\ \mathbf{m} = (D\mathbf{n} + C\mathbf{k})\ , \qquad & m_0 = (t n_0 + s k_0)\ . \end{aligned} \tag{10.1}$$

The multiplication law takes the form

$$\begin{aligned} k_0'' &= k_0'\, k_0 + \mathbf{k}' \cdot \mathbf{k} + n_0'\, (\alpha k_0 + \beta n_0) + \mathbf{n}'\, (A\mathbf{k} + B\mathbf{n})\ , \\ m_0'' &= (t n_0' + s k_0')\, (t n_0 + s k_0) + (D\mathbf{n}' + C\mathbf{k}')\, (D\mathbf{n} + C\mathbf{k}) \\ &\quad + (\alpha k_0' + \beta n_0') n_0 + (A\mathbf{k}' + B\mathbf{n}')\, \mathbf{n}\ , \\ n_0'' &= k_0'\, n_0 + \mathbf{k}' \cdot \mathbf{n} + n_0'\, (t n_0 + s k_0) + \mathbf{n}'\, (D\mathbf{n} + C\mathbf{k})\ , \\ l_0'' &= (\alpha k_0' + \beta n_0')\, k_0 + (A\mathbf{k}' + B\mathbf{n}')\, \mathbf{k} \\ &\quad + (t n_0' + s k_0')(\alpha k_0 + \beta n_0) + (D\mathbf{n}' + C\mathbf{k}')\, (A\mathbf{k} + B\mathbf{n})\ , \end{aligned}$$

$$\begin{aligned} \mathbf{k}'' &= k_0'\, \mathbf{k} + \mathbf{k}'\, k_0 + i\, \mathbf{k}' \times \mathbf{k} + n_0'\, (A\mathbf{k} \\ &\quad + B\mathbf{n}) + \mathbf{n}'\, (\alpha k_0 + \beta n_0) + i\, \mathbf{n}' \times (A\mathbf{k} + B\mathbf{n})\ , \\ \mathbf{m}'' &= (t n_0' + s k_0')(D\mathbf{n} + C\mathbf{k}) + (D\mathbf{n}' + C\mathbf{k}')\, (t n_0 + s k_0) \\ &\quad + i\, (D\mathbf{n}' + C\mathbf{k}') \times (D\mathbf{n} + C\mathbf{k}) \\ &\quad + (\alpha k_0' + \beta n_0')\, \mathbf{n} + (A\mathbf{k}' + B\mathbf{n}')\, n_0 + i\, (A\mathbf{k}' + B\mathbf{n}') \times \mathbf{n}\ , \\ \mathbf{n}'' &= k_0'\, \mathbf{n} + \mathbf{k}'\, n_0 + i\, \mathbf{k}' \times \mathbf{n} + n_0'\, (D\mathbf{n} + C\mathbf{k}) \\ &\quad + \mathbf{n}'\, (t n_0 + s k_0) + i\, \mathbf{n}' \times (D\mathbf{n} + C\mathbf{k})\ , \\ \mathbf{l}'' &= (\alpha k_0' + \beta n_0')\, \mathbf{k} + (A\mathbf{k}' + B\mathbf{n}')\, k_0 + i\, (A\mathbf{k}' + B\mathbf{n}') \times \mathbf{k} \\ &\quad + (t n_0' + s k_0')(A\mathbf{k} + B\mathbf{n}) + (D\mathbf{n}' + C\mathbf{k}')(\alpha k_0 + \beta n_0) \\ &\quad + i\, (D\mathbf{n}' + C\mathbf{k}') \times (A\mathbf{k} + B\mathbf{n})\ . \end{aligned} \tag{10.2}$$

The equation

$$\mathbf{l}'' = (A\mathbf{k}'' + B\mathbf{n}'') \tag{10.3}$$

is equivalent to

$$\begin{aligned}
&(\alpha k_0' + \beta n_0')\mathbf{k} + (A\mathbf{k}' + B\mathbf{n}')k_0 + i(A\mathbf{k}' + B\mathbf{n}') \times \mathbf{k} \\
&+(tn_0' + sk_0')(A\mathbf{k} + B\mathbf{n}) + (D\mathbf{n}' + C\mathbf{k}')(\alpha k_0 + \beta n_0) + i(D\mathbf{n}' + C\mathbf{k}') \times (A\mathbf{k} + B\mathbf{n}) \\
&= A[k_0'\mathbf{k} + \mathbf{k}'k_0 + i\mathbf{k}' \times \mathbf{k} + n_0'(A\mathbf{k} + B\mathbf{n}) + \mathbf{n}'(\alpha k_0 + \beta n_0) + i\mathbf{n}' \times (A\mathbf{k} + B\mathbf{n})] \\
&+B[k_0'\ \mathbf{n} + \mathbf{k}'n_0 + i\mathbf{k}' \times \mathbf{n} + n_0'(D\mathbf{n} + C\mathbf{k}) + \mathbf{n}'(tn_0 + sk_0) + i\mathbf{n}' \times (D\mathbf{n} + C\mathbf{k})];
\end{aligned}$$

this leads to

$$\begin{aligned}
k_0'\mathbf{k}: &\quad \alpha + sA = A\ , \quad k_0\mathbf{k}': \quad A + \alpha C = A\ , \\
n_0\mathbf{n}': &\quad \beta D = \beta A + tB\ , \quad n_0'\mathbf{n}: \quad tB = AB + BD\ , \\
k_0'\mathbf{n}: &\quad sB = B\ , \quad k_0\mathbf{n}': \quad B + \alpha D = \alpha A + sB\ , \\
n_0'\mathbf{k}: &\quad \beta + tA = A^2 + BC\ , \quad n_0\mathbf{k}': \quad \beta C = B\ , \\
\mathbf{k}' \times \mathbf{k}: &\quad A + AC = A\ , \quad \mathbf{n}' \times \mathbf{n}: \quad BD = AB + BD\ , \\
\mathbf{n}' \times \mathbf{k}: &\quad B + AD = A^2 + BC\ , \quad \mathbf{k}' \times \mathbf{n}: \quad CB = B\ .
\end{aligned}$$

The equation

$$\mathbf{m}'' = (D\mathbf{n}'' + C\mathbf{k}'') \tag{10.4}$$

takes the form

$$\begin{aligned}
&(tn_0' + sk_0')(D\mathbf{n} + C\mathbf{k}) + (D\mathbf{n}' + C\mathbf{k}')(tn_0 + sk_0) + i(D\mathbf{n}' + C\mathbf{k}') \times (D\mathbf{n} + C\mathbf{k}) \\
&+(\alpha k_0' + \beta n_0')\mathbf{n} + (A\mathbf{k}' + B\mathbf{n}')n_0 + i(A\mathbf{k}' + B\mathbf{n}') \times \mathbf{n} \\
&= D[k_0'\mathbf{n} + \mathbf{k}'n_0 + i\mathbf{k}' \times \mathbf{n} + n_0'(D\mathbf{n} + C\mathbf{k}) + \mathbf{n}'(tn_0 + sk_0) + i\mathbf{n}' \times (D\mathbf{n} + C\mathbf{k})] \\
&+C[k_0'\mathbf{k} + \mathbf{k}'k_0 + i\mathbf{k}' \times \mathbf{k} + n_0'(A\mathbf{k} + B\mathbf{n}) + \mathbf{n}'(\alpha k_0 + \beta n_0) + i\mathbf{n}' \times (A\mathbf{k} + B\mathbf{n})];
\end{aligned}$$

whence it follows

$$\begin{aligned}
k_0'\mathbf{k}: &\quad sC = C\ , \quad k_0\mathbf{k}': \quad sC = C\ , \\
n_0\mathbf{n}': &\quad tD + B = tD + \beta C\ , \quad n_0'\mathbf{n}: \quad tD + \beta = D^2 + CB\ , \\
k_0'\mathbf{n}: &\quad sD + \alpha = D\ , \quad k_0\mathbf{n}': \quad sD = sD + \alpha C\ , \\
n_0'\mathbf{k}: &\quad tC = DC + CA\ , \quad n_0\mathbf{k}': \quad tC + A = D\ , \\
\mathbf{k}' \times \mathbf{k}: &\quad C^2 = C\ , \quad \mathbf{n}' \times \mathbf{n}: \quad D^2 + B = D^2 + CB\ , \\
\mathbf{n}' \times \mathbf{k}: &\quad DC = DC + CA\ , \quad \mathbf{k}' \times \mathbf{n}: \quad CD + A = D\ .
\end{aligned}$$

The equation

$$l_0'' = (\alpha k_0'' + \beta n_0'') \tag{10.5}$$

gives

$$\begin{aligned}
&(\alpha k_0' + \beta n_0')\ k_0 + (A\mathbf{k}' + B\mathbf{n}')\ \mathbf{k} \\
&+(tn_0' + sk_0')(\alpha k_0 + \beta n_0) + (D\mathbf{n}' + C\mathbf{k}')\ (A\mathbf{k} + B\mathbf{n}) \\
&= \alpha\ [\ k_0'\ k_0 + \mathbf{k}' \cdot \mathbf{k} + n_0'\ (\alpha k_0 + \beta n_0) + \mathbf{n}'\ (A\mathbf{k} + B\mathbf{n})\ ] \\
&+\beta\ [\ (tn_0' + sk_0')\ (tn_0 + sk_0) + (D\mathbf{n}' + C\mathbf{k}')\ (D\mathbf{n} + C\mathbf{k})\ ]\ ,
\end{aligned}$$

which leads to

$$\begin{aligned}
k_0'k_0: &\quad \alpha + \alpha s = \alpha + \beta s^2\ , \quad n_0'n_0: \quad t\beta = \alpha\beta + \beta t^2\ , \\
k_0'n_0: &\quad s\beta = \beta st\ , \quad k_0n_0': \quad \beta + t\alpha = \alpha^2 + \beta ts\ , \\
\mathbf{k}' \cdot \mathbf{k}: &\quad A + CA = \alpha + \beta C^2\ , \quad \mathbf{n}' \cdot \mathbf{n}: \quad DB = \alpha B + \beta D^2\ ,
\end{aligned}$$

$$\mathbf{k}'\cdot\mathbf{n}: \qquad CB=\beta CD\ , \quad \mathbf{k}\cdot\mathbf{n}': \qquad B+AD=\alpha A+\beta CD\ .$$

The equation

$$m_0''=tn_0''+sk_0'' \tag{10.6}$$

is equivalent to

$$\begin{aligned}(tn_0'+sk_0')\,(tn_0+sk_0)+(D\mathbf{n}'+C\mathbf{k}')\,(D\mathbf{n}+C\mathbf{k})\\ +(\alpha k_0'+\beta n_0')n_0+(A\mathbf{k}'+B\mathbf{n}')\,\mathbf{n}\\ =t\,[\,k_0'\,n_0+\mathbf{k}'\cdot\mathbf{n}+n_0'\,(tn_0+sk_0)+\mathbf{n}'\,(D\mathbf{n}+C\mathbf{k})\\ +s[\,k_0'\,k_0+\mathbf{k}'\cdot\mathbf{k}+n_0'\,(\alpha k_0+\beta n_0)+\mathbf{n}'\,(A\mathbf{k}+B\mathbf{n})\,]\ ;\end{aligned}$$

whence it follows

$$\begin{aligned} k_0'k_0: &\qquad s^2=s\ , \quad n_0'n_0: &&\qquad t^2+\beta=t^2+s\beta\ ,\\ k_0'n_0: &\qquad st+\alpha=t\ , \quad k_0n_0': &&\qquad ts=st+\alpha s\ ,\\ \mathbf{k}'\cdot\mathbf{k}: &\qquad C^2=s\ , \quad \mathbf{n}'\cdot\mathbf{n}: &&\qquad D^2+B=tD+sB\ ,\\ \mathbf{k}'\cdot\mathbf{n}: &\qquad CD+A=t\ , \quad \mathbf{n}'\cdot\mathbf{k}: &&\qquad DC=tC+sA\ .\end{aligned}$$

We collect results together, and yield

$$\alpha+sA=A\ ,\ \alpha C=0\ ,\ \beta D=\beta A+tB\ ,\ tB=AB+BD\ ,\ sB=B\ ,$$

$$B+\alpha D=\alpha A+sB\ ,\ \beta+tA=A^2+BC\ ,\ \beta C=B\ ,\ AC=0\ ,\ 0=AB\ ,$$

$$B+AD=A^2+BC\ ,\ CB=B\ ;\ sC=C\ ,\ sC=C\ ,$$

$$B=\beta C\ ,\ tD+\beta=D^2+CB\ ,\ sD+\alpha=D\ ,\ 0=\alpha C\ ,$$

$$tC=DC+CA,\ tC+A=D,\ C(C-1)=0,$$

$$B=CB,\ 0=CA,\ CD+A=D,\ \alpha s=\beta s^2,$$

$$t\beta=\alpha\beta+\beta t^2,\ s\beta=\beta st,\ \beta+t\alpha=\alpha^2+\beta ts,\ A+CA=\alpha+\beta C^2,$$

$$DB=\alpha B+\beta D^2,\ CB=\beta CD,\ B+AD=\alpha A+\beta CD;$$

$$s(s-1)=0\ ,\ \beta=s\beta\ ,\ st+\alpha=t\ ,\ 0=\alpha s\ ,\ C^2=s\ ,$$

$$D^2+B=tD+sB\ ,\ CD+A=t\ ,\ DC=tC+sA\ .$$

We see that for s, C there are possible only the following values

$$C=s=0\ , \qquad C=s=1\ .$$

If $C=s=0$, we get only one

solution $(KN-1)$,

$$\begin{aligned}&\alpha=A\ , \quad B=\beta=0\ , \quad C=s=0\ , \quad D=t=A\ ,\\ &\mathbf{l}=A\mathbf{k}\ , \quad l_0=Ak_0\ , \quad \mathbf{m}=A\mathbf{n}\ ,\ m_0=An_0\ ,\end{aligned} \tag{10.7}$$

$$G=\begin{pmatrix} K & N\\ AK & AN\end{pmatrix},\ G'G=\begin{pmatrix} (K'K+AN'K) & (K'N+AN'N)\\ A(K'K+AN'K) & A(K'N+AN'N)\end{pmatrix}.$$

In the case $C=s=+1$, we also get only one

solution $(KN-2)$,

$$\begin{aligned}&A=\alpha=0\ , \quad B=\beta=0\ , \quad C=s=+1\ , \quad D=t=0\ ,\\ &\mathbf{l}=0\ , \quad l_0=0\ , \quad \mathbf{m}=\mathbf{k}\ , \quad m_0=k_0\ ,\end{aligned} \tag{10.8}$$

$$G=\begin{pmatrix} K & N\\ 0 & K\end{pmatrix},\ G'G=\begin{pmatrix} K'K & (K'N+N'K)\\ 0 & K'K\end{pmatrix}.$$

11 Two Independent Vectors: Variant II(m,l)

Let us examine **variant II(m,l)**:

$$\begin{aligned} \mathbf{n} &= A\mathbf{m} + B\mathbf{l}\,, \qquad & l_0 &= \alpha m_0 + \beta l_0\,, \\ \mathbf{k} &= D\mathbf{l} + C\mathbf{m}\,, \qquad & k_0 &= t l_0 + s m_0\,. \end{aligned} \tag{11.1}$$

Here there arise only

solution $(ML-1)$,

$$G = SG'S^{-1} = \begin{pmatrix} AN' & AK' \\ N' & K' \end{pmatrix} \equiv \begin{pmatrix} AL & AM \\ L & M \end{pmatrix}; \tag{11.2}$$

solution $(ML-2)$,

$$G = SG'S^{-1} = \begin{pmatrix} K' & 0 \\ N' & K' \end{pmatrix} \equiv \begin{pmatrix} M & 0 \\ L & M \end{pmatrix}. \tag{11.3}$$

12 Three Independent Vectors: Variants I(k,m,n) and I(m,k,l)

Now consider the case of three independent vectors

$$\mathbf{l} = A\mathbf{k} + B\mathbf{m} + C\mathbf{n}\,, \qquad l_0 = \alpha k_0 + \beta m_0 + s n_0\,. \tag{12.1}$$

First, let us examine the simpler second restriction in (12.1). The formulas for k_0'', m_0'', n_0'', l_0'' take the form

$$\begin{aligned} k_0'' &= k_0' k_0 + \mathbf{k}'\cdot\mathbf{k} + n_0'(\alpha k_0 + \beta m_0 + s n_0) + \mathbf{n}'(A\mathbf{k} + B\mathbf{m} + C\mathbf{n})\,, \\ m_0'' &= m_0' m_0 + \mathbf{m}'\cdot\mathbf{m} + (\alpha k_0' + \beta m_0' + s n_0')n_0 + (A\mathbf{k}' + B\mathbf{m}' + C\mathbf{n}')\mathbf{n}\,, \end{aligned}$$

$$\begin{aligned} n_0'' &= k_0'\, n_0 + \mathbf{k}'\cdot\mathbf{n} + n_0'\, m_0 + \mathbf{n}'\cdot\mathbf{m}\,, \\ l_0'' &= (\alpha k_0' + \beta m_0' + s n_0')k_0 + (A\mathbf{k} + B\mathbf{m} + C\mathbf{n})\mathbf{k} \\ &+ m_0'(\alpha k_0 + \beta m_0 + s n_0) + \mathbf{m}'(A\mathbf{k} + B\mathbf{m} + C\mathbf{n})\,. \end{aligned}$$

We set

$$l_0'' = \alpha k_0'' + \beta m_0'' + s n_0'' \tag{12.2}$$

or, equivalently,

$$\begin{aligned} &(\alpha k_0' + \beta m_0' + s n_0')k_0 + (A\mathbf{k}' + B\mathbf{m}' + C\mathbf{n}')\mathbf{k} \\ &+ m_0'(\alpha k_0 + \beta m_0 + s n_0) + \mathbf{m}'(A\mathbf{k} + B\mathbf{m} + C\mathbf{n}) \\ &= \alpha\,[\, k_0' k_0 + \mathbf{k}'\cdot\mathbf{k} + n_0'(\alpha k_0 + \beta m_0 + s n_0) + \mathbf{n}'(A\mathbf{k} + B\mathbf{m} + C\mathbf{n})\,] \\ &+ \beta[\, m_0' m_0 + \mathbf{m}'\cdot\mathbf{m} + (\alpha k_0' + \beta m_0' + s n_0')n_0 + (A\mathbf{k}' + B\mathbf{m}' + C\mathbf{n}')\mathbf{n}\,] \\ &+ s\,[\, k_0'\, n_0 + \mathbf{k}'\cdot\mathbf{n} + n_0'\, m_0 + \mathbf{n}'\cdot\mathbf{m}\,]\,, \end{aligned}$$

which results in

$$\begin{aligned} k_0' k_0 &:\ \alpha = \alpha\,, & m_0' m_0 &:\ \beta = \beta;\,, \\ m_0' k_0 &:\ \beta + \alpha = 0\,, & n_0' n_0 &:\ 0 = (\alpha+\beta)s\,, \\ n_0' k_0 &:\ s = \alpha^2\,, & m_0' n_0 &:\ s = \beta^2\,, \\ n_0' m_0 &:\ 0 = \beta(\alpha + s)\,, & k_0' n_0 &:\ 0 = \beta(\alpha + s)\,, \\ \mathbf{k}'\cdot\mathbf{k} &:\ A = \alpha\,, & \mathbf{m}'\cdot\mathbf{m} &:\ B = \beta\,, \\ \mathbf{m}'\cdot\mathbf{k} &:\ B + A = 0\,, & \mathbf{n}'\cdot\mathbf{n} &:\ 0 = (\alpha+\beta)C\,, \\ \mathbf{n}'\cdot\mathbf{k} &:\ C = \alpha A & \mathbf{m}'\cdot\mathbf{n} &:\ C = \beta B\,, \\ \mathbf{n}'\cdot\mathbf{m} &:\ 0 = \alpha B + \beta s\,, & \mathbf{k}'\cdot\mathbf{n} &:\ 0 = \beta A + \beta s\,. \end{aligned} \tag{12.3}$$

Let us write down only the independent equations:

$$\alpha = A, \qquad B = \beta = -A, \qquad C = s = A^2, \qquad A(A+s) = 0\ . \tag{12.4}$$

The equations (12.4) allow two types of solutions:

$$(I), \qquad A = \alpha = 0\ , \qquad B = \beta = 0\ ,\ C = s = 0\ ; \tag{12.5}$$

$$(II), \qquad A = \alpha = -1, \qquad B = \beta = +1\ , \qquad C = s = +1\ . \tag{12.6}$$

It remains to verify that these solutions satisfy the equation (see (12.1))

$$\mathbf{l}'' = A\mathbf{k}'' + B\mathbf{m}'' + C\mathbf{n}''\ . \tag{12.7}$$

Evidently, we should check only the type II – it takes the form

$$\mathbf{l}'' = -\mathbf{k}'' + \mathbf{m}'' + \mathbf{n}''$$

and further reduces to

$$\begin{aligned}
&(-k_0' + m_0' + n_0')\ \mathbf{k} + (-\mathbf{k}' + \mathbf{m}' + \mathbf{n}')\ k_0 \\
&+i\ (-\mathbf{k}' + \mathbf{m}' + \mathbf{n}') \times \mathbf{k} + m_0'\ (-\mathbf{k} + \mathbf{m} + \mathbf{n}) \\
&+\mathbf{m}'(-k_0 + m_0 + n_0) + i\ \mathbf{m}' \times (-\mathbf{k} + \mathbf{m} + \mathbf{n}) \\
&= -[\ k_0'\ \mathbf{k} + \mathbf{k}'\ k_0 + i\ \mathbf{k}' \times \mathbf{k} + n_0'\ (-\mathbf{k} + \mathbf{m} + \mathbf{n}) \\
&+\mathbf{n}'\ (-k_0 + m_0 + n_0) + i\ \mathbf{n}' \times (-\mathbf{k} + \mathbf{m} + \mathbf{n})\ ] \\
&+[\ m_0'\ \mathbf{m} + \mathbf{m}'\ m_0 + i\ \mathbf{m}' \times \mathbf{m} + (-k_0' + m_0' + n_0')\ \mathbf{n} + \\
&(-\mathbf{k}' + \mathbf{m}' + \mathbf{n}')\ n_0 + i\ (-\mathbf{k}' + \mathbf{m}' + \mathbf{n}') \times \mathbf{n}\ ] \\
&+[\ k_0'\ \mathbf{n} + \mathbf{k}'\ n_0 + i\ \mathbf{k}' \times \mathbf{n} + n_0'\ \mathbf{m} + \mathbf{n}'\ m_0 + i\ \mathbf{n}' \times \mathbf{m}\ ]\ ,
\end{aligned}$$

which turns to be an identity.

Thus, we have constructed two solutions:

solution $(KMN-1)$,

$$G = \begin{pmatrix} K & N \\ 0 & M \end{pmatrix},\ G'G = \begin{pmatrix} K'K & (K'N + N'M) \\ 0 & M'M \end{pmatrix}; \tag{12.8}$$

solution $(KMN-2)$,

$$\begin{aligned}
&\mathbf{l} = -\mathbf{k} + \mathbf{m} + \mathbf{n}\ ,\ l_0 = -k_0 + m_0 + n_0\ , \\
&G = \begin{pmatrix} K & N \\ -K + M + N & M \end{pmatrix};
\end{aligned} \tag{12.9}$$

These are sets of degenerate matrices of rank 2. It remains to verify the multiplication law for the case (12.9):

$$\begin{aligned}
&\begin{pmatrix} K' & N' \\ -K' + M' + N' & M' \end{pmatrix} \begin{pmatrix} K & N \\ -K + M + N & M \end{pmatrix} \\
&= \begin{pmatrix} (K'K - N'K + N'M + N'N) & K'N + N'M \\ (-K'K + N'K + M'M + M'N) & (-K'N + M'N + N'N + M'M) \end{pmatrix}\ .
\end{aligned} \tag{12.10}$$

With the notations

$$\begin{aligned}
K'' &= K'K - N'K + N'M + N'N\ , \\
M'' &= -K'N + M'N + N'N + M'M\ , \quad N'' = K'N + N'M\ ,
\end{aligned}$$

we obtain

$$-K'' + M'' + N'' = -K'K + N'K - N'M \ ,$$
$$-N'N - K'N + M'N + N'N + M'M + K'N + N'M = -K'K + N'K + M'N + M'M \ ;$$

Thus, the set (12.9) has semi-group structure

$$G'G = \begin{pmatrix} K'' & N'' \\ -K'' + M'' + N'' & M' \end{pmatrix} . \tag{12.11}$$

For the variant **II(k,m,l)** one has the similar solutions:

solution $(KML-1)$,

$$G = \begin{pmatrix} K & 0 \\ L & M \end{pmatrix} ; \tag{12.12}$$

solution $(KML-2)$,

$$G = \begin{pmatrix} K & -M+K+L \\ L & M \end{pmatrix} . \tag{12.13}$$

13 Three Independent Vectors: Variants I(n,l,k) and I(n,l,m)

Let us consider the variant $I(n,l,k)$

$$\mathbf{m} = A\mathbf{n} + B\mathbf{l} + C\mathbf{k} \ , \qquad m_0 = \alpha n_0 + \beta l_0 + s k_0 \ . \tag{13.1}$$

First, we turn to the simplest second restriction in (13.1); this gives

$$\begin{aligned} k_0'' &= k_0' \ k_0 + \mathbf{k}' \cdot \mathbf{k} + n_0' \ l_0 + \mathbf{n}' \cdot \mathbf{l} \ , \\ m_0'' &= (\alpha n_0' + \beta l_0' + s k_0')(\alpha n_0 + \beta l_0 + s k_0) \\ &+ (A\mathbf{n}' + B\mathbf{l}' + C\mathbf{k}')(A\mathbf{n} + B\mathbf{l} + C\mathbf{k}) + l_0' \ n_0 + \mathbf{l}' \cdot \mathbf{n} \ , \\ n_0'' &= k_0' \ n_0 + \mathbf{k}' \cdot \mathbf{n} + n_0'(\alpha n_0 + \beta l_0 + s k_0) + \mathbf{n}' \ (A\mathbf{n} + B\mathbf{l} + C\mathbf{k}) \ , \\ l_0'' &= l_0' \ k_0 + \mathbf{l}' \cdot \mathbf{k} + (\alpha n_0' + \beta l_0' + s k_0') l_0 + (A\mathbf{n}' + B\mathbf{l}' + C\mathbf{k}')\mathbf{l} \ . \end{aligned} \tag{13.2}$$

By requiring

$$m_0'' = \alpha n_0'' + \beta l_0'' + s k_0'' \ ,$$

or

$$\begin{aligned} &(\alpha n_0' + \beta l_0' + s k_0')(\alpha n_0 + \beta l_0 + s k_0) \\ &+ (A\mathbf{n}' + B\mathbf{l}' + C\mathbf{k}')(A\mathbf{n} + B\mathbf{l} + C\mathbf{k}) + l_0' \ n_0 + \mathbf{l}' \cdot \mathbf{n} \\ &= \alpha \ [\ k_0' \ n_0 + \mathbf{k}' \cdot \mathbf{n} + n_0'(\alpha n_0 + \beta l_0 + s k_0) + \mathbf{n}' \ (A\mathbf{n} + B\mathbf{l} + C\mathbf{k}) \ ] \\ &+ \beta \ [\ l_0' \ k_0 + \mathbf{l}' \cdot \mathbf{k} + (\alpha n_0' + \beta l_0' + s k_0') l_0 + (A\mathbf{n}' + B\mathbf{l}' + C\mathbf{k}')\mathbf{l} \ ] \\ &+ s \ [\ k_0' \ k_0 + \mathbf{k}' \cdot \mathbf{k} + n_0' \ l_0 + \mathbf{n}' \cdot \mathbf{l} \ ] \ , \end{aligned}$$

we obtain the equations

$$\begin{aligned} n_0' n_0 : \ \alpha^2 = \alpha^2 \ , \qquad l_0' l_0 : \ \beta^2 = \beta^2 \ , \qquad k_0' k_0 : \ s^2 = s \ , \\ n_0' l_0 : \ \alpha\beta = \alpha\beta + \alpha\beta + s \ , \qquad n_0' k_0 : \ \alpha s = \alpha s \ , \\ l_0' n_0 : \ \beta\alpha + 1 = 0 \ , \qquad l_0' k_0 : \ \beta s = \beta \ , \\ k_0' n_0 : \ s\alpha = \alpha \ , \qquad k_0' l_0 : \ s\beta = \beta s \ , \end{aligned} \tag{13.3}$$

$$\mathbf{n}'\cdot\mathbf{n}:\ A^2=\alpha A,\qquad \mathbf{l}'\cdot\mathbf{l}:\ B^2=\beta B\ ,\qquad \mathbf{k}'\cdot\mathbf{k}:\ C^2=s\ ,$$
$$\mathbf{n}'\cdot\mathbf{l}:\ AB=\alpha B+\beta A+s\ ,\qquad \mathbf{n}'\cdot\mathbf{k}:\ AC=\alpha C\ ,$$
$$\mathbf{l}'\cdot\mathbf{n}:\ BA+1=0\ ,\qquad \mathbf{l}'\cdot\mathbf{k}:\ BC=\beta\ ,$$
$$\mathbf{k}'\cdot\mathbf{n}:\ CA=\alpha\ ,\qquad \mathbf{k}'\cdot\mathbf{l}:\ CB=\beta C\ . \tag{13.4}$$

Let us write down only the independent equations:

$$\alpha\beta=-1,\qquad s=-\alpha\beta=+1,\qquad C^2=s=1\ ,$$
$$A(A-\alpha)=0,\qquad C(A-\alpha)=0,\qquad B(B-\beta)=0,$$
$$AB=\alpha B+\beta A+1\ ,\qquad AC=\alpha C\ ,$$
$$BA+1=0\ ,\quad BC=\beta\ ,\quad CA=\alpha\ ,\quad CB=\beta C\ ;$$

they provide to us only with

solution $(NLK-1)$,

$$\alpha=A,\qquad B=\beta=-\frac{1}{A}\ ,\qquad C=s=+1,$$
$$\mathbf{m}=\mathbf{k}+(A\,\mathbf{n}-A^{-1}\,\mathbf{l})\ ,\qquad m_0=k_0+(A\,n_0-A^{-1}l_0)\ ; \tag{13.5}$$

where the corresponding matrices have the structure

$$G=\begin{pmatrix} K & N \\ L & (K+AN-A^{-1}L) \end{pmatrix}. \tag{13.6}$$

This is a set of degenerate matrices of rank 2. We verify now the multiplication law:

$$G'G=\begin{pmatrix} K' & N' \\ L' & (K'+AN'-A^{-1}L') \end{pmatrix}\begin{pmatrix} K & N \\ L & (K+AN-A^{-1}L) \end{pmatrix}$$
$$\begin{pmatrix} (K'K+N'L) & K'N+N'(K+AN-A^{-1}L) \\ L'K+(K'+AN'-A^{-1}L')L & L'N+(K'+AN'-A^{-1}L')(K+AN-A^{-1}L) \end{pmatrix}.$$

The block (22) in the matrix (13.7) is

$$(22)=L'N+(K'+AN'-A^{-1}L')\,(K+AN-A^{-1}L)$$
$$=(K'K-N'L)+A\,(K'N+N'K)-A^{-1}\,(K'L+L'K)+A^2N'N+A^{-2}L'L\ .$$

With the notations

$$K''=K'K+N'L\ ,$$
$$N''=K'N+N'(K+AN-A^{-1}L)\ ,$$
$$L''=L'K+(K'+AN'-A^{-1}L')L\ ,$$

we get

$$K''+AN''-A^{-1}L''=K'K+N'L+A\,[\,K'N+N'(K+AN-A^{-1}L)\,]$$
$$-A^{-1}\,[\,L'K+(K'+AN'-A^{-1}L')L\,]=(K'K-N'L)+A\,(K'N+N'K)$$
$$-A^{-1}\,(L'K+K'L)+(A^2N'N+A^{-2}L'L)\ ,$$

and we obtain the claimed relation

$$(22)=K''+AN''-A^{-1}L''\ .$$

Therefore we have

$$G'G=\begin{pmatrix} K'' & N'' \\ L'' & (K''+AN''-A^{-1}L'') \end{pmatrix}=G''\ .$$

There exists one solution for the variant $I(n,l,m)$:
solution $(NLM-1)$,

$$G = \begin{pmatrix} (M + AL - A^{-1}N) & N \\ L & M \end{pmatrix} . \tag{13.7}$$

14 Degenerate Matrices of Rank 3

Let us consider singular Mueller matrices of rank 3. Given the explicit form of the matrix G, it is easy to understand that there are 16 simple ways to get the semigroups of rank 3. For this, it suffices to have vanishing some i-line or some j-column in the original 4-dimensional matrix,

$$\begin{array}{cccc} (00) , & (01) , & (02) , & (03) , \\ (10) , & (11) , & (12) , & (13) , \\ (20) , & (21) , & (22) , & (23) , \\ (30) , & (31) , & (32) , & (33) . \end{array} \tag{14.1}$$

The compatibility of the law of multiplication with this constraint is obvious.
All the 16 possibilities are listed below.

Variant (00)

$$G = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 2k_0 & 2n_1 & 2n_0 \\ 0 & 2l_1 & m_0 + m_3 & m_1 - im_2 \\ 0 & 2l_0 & m_1 + im_2 & m_0 - m_3 \end{pmatrix},$$

$$k_1 = 0, \qquad k_2 = 0, \qquad k_0 = -k_3 ,$$

$$n_0 = -n_3, \qquad l_0 = -l_3, \qquad +in_2 = n_1, \qquad -il_2 = l_1 .$$

Variant (01)

$$G = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 2k_1 & 0 & 2n_1 & 2n_0 \\ 2l_0 & 0 & m_0 + m_3 & m_1 - im_2 \\ 2l_1 & 0 & m_1 + im_2 & m_0 - m_3 \end{pmatrix},$$

$$k_0 = 0, \qquad k_3 = 0, \qquad k_1 = ik_2 ,$$

$$l_1 = il_2, \qquad l_0 = l_3, \qquad n_0 = -n_3, \qquad n_1 = in_2 .$$

Variant (02)

$$G = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 2k_1 & 2k_0 & 0 & 2n_0 \\ l_0 + l_3 & l_1 - il_2 & 0 & 2m_1 \\ l_1 + il_2 & l_0 - l_3 & 0 & 2m_0 \end{pmatrix},$$

$$n_1 = 0, \qquad n_2 = 0, \qquad n_0 = -n_3 ,$$

$$m_0 = -m_3, \qquad m_1 = -im_2, \qquad k_0 = -k_3, \qquad k_1 = ik_2 .$$

Variant (03)

$$G = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 2k_1 & 2k_0 & 2n_1 & 0 \\ l_0 + l_3 & l_1 - il_2 & 2m_0 & 0 \\ l_1 + il_2 & l_0 - l_3 & 2m_1 & 0 \end{pmatrix},$$

$$n_0 = 0, \qquad n_3 = 0, \qquad n_1 = in_2 ,$$

$$m_0 = m_3, \qquad m_1 = im_2, \qquad k_0 = -k_3, \qquad k_1 = ik_2 .$$

Variant (10)

$$G = \begin{pmatrix} 0 & 2k_1 & 2n_0 & 2n_1 \\ 0 & 0 & 0 & 0 \\ 0 & 2l_1 & m_0+m_3 & m_1-im_2 \\ 0 & 2l_0 & m_1+im_2 & m_0-m_3 \end{pmatrix},$$

$$k_0 = 0, \qquad k_3 = 0, \qquad k_1 = -ik_2\ ,$$

$$l_0 = -l_3, \qquad l_1 = -il_2, \qquad n_1 = -in_2, \qquad n_0 = n_3\ .$$

Variant (11)

$$G = \begin{pmatrix} 2k_0 & 0 & 2n_0 & 2n_1 \\ 0 & 0 & 0 & 0 \\ 2l_0 & 0 & m_0+m_3 & m_1-im_2 \\ 2l_1 & 0 & m_1+im_2 & m_0-m_3 \end{pmatrix},$$

$$k_1 = 0, \qquad k_2 = 0, \qquad k_0 = k_3\ ,$$

$$l_0 = l_3, \qquad l_1 = il_2, \qquad n_1 = -in_2, \qquad n_0 = n_3\ .$$

Variant (12)

$$G = \begin{pmatrix} 2k_0 & 2k_1 & 0 & 2n_1 \\ 0 & 0 & 0 & 0 \\ l_0+l_3 & l_1-il_2 & 0 & 2m_1 \\ l_1+il_2 & l_0-l_3 & 0 & 2m_0 \end{pmatrix},$$

$$n_0 = 0, \qquad n_3 = 0, \qquad n_1 = -in_2\ ,$$

$$m_0 = -m_3, \qquad m_1 = -im_2, \qquad k_1 = -ik_2, \qquad k_0 = k_3\ .$$

Variant (13)

$$G = \begin{pmatrix} 2k_0 & 2k_1 & 2n_0 & 0 \\ 0 & 0 & 0 & 0 \\ l_0+l_3 & l_1-il_2 & 2m_0 & 0 \\ l_1+il_2 & l_0-l_3 & 2m_1 & 0 \end{pmatrix},$$

$$n_1 = 0, \qquad n_2 = 0, \qquad n_0 = n_3\ ,$$

$$m_0 = m_3, \qquad m_1 = im_2, \qquad k_1 = -ik_2, \qquad k_0 = k_3\ .$$

Variant (20)

$$G = \begin{pmatrix} 0 & 2k_1 & n_0+n_3 & n_1-in_2 \\ 0 & 2k_0 & n_1+in_2 & n_0-n_3 \\ 0 & 0 & 0 & 0 \\ 0 & 2l_0 & 2m_1 & 2m_0 \end{pmatrix},$$

$$l_1 = 0, \qquad l_2 = 0, \qquad l_0 = -l_3\ ,$$

$$m_0 = -m_3, \qquad m_1 = im_2, \qquad k_1 = -ik_2, \qquad k_0 = -k_3\ .$$

Variant (21)

$$G = \begin{pmatrix} 2k_0 & 0 & n_0+n_3 & n_1-in_2 \\ 2k_1 & 0 & n_1+in_2 & n_0-n_3 \\ 0 & 0 & 0 & 0 \\ 2l_1 & 0 & 2m_1 & 2m_0 \end{pmatrix},$$

$$l_0 = 0, \qquad l_3 = 0, \qquad l_1 = il_2\ ,$$

$$m_0 = -m_3, \qquad m_1 = im_2, \qquad k_1 = ik_2, \qquad k_0 = k_3\ .$$

Variant (22)

$$G = \begin{pmatrix} k_0+k_3 & k_1-ik_2 & 0 & 2n_1 \\ k_1+ik_2 & k_0-k_3 & 0 & 2n_0 \\ 0 & 0 & 0 & 0 \\ 2l_1 & 2l_0 & 0 & 2m_0 \end{pmatrix},$$

$$m_1 = 0, \qquad m_2 = 0, \qquad m_0 = -m_3\ ,$$

$$n_0 = -n_3, \qquad n_1 = -in_2, \qquad l_1 = il_2, \qquad l_0 = -l_3\ .$$

Variant (23)

$$G = \begin{pmatrix} k_0+k_3 & k_1-ik_2 & 2n_0 & 0 \\ k_1+ik_2 & k_0-k_3 & 2n_1 & 0 \\ 0 & 0 & 0 & 0 \\ 2l_1 & 2l_0 & 2m_1 & 0 \end{pmatrix}$$

$$m_0 = 0, \qquad m_3 = 0, \qquad m_1 = im_2\ ,$$

$$n_0 = n_3, \qquad n_1 = in_2, \qquad l_1 = il_2, \qquad l_0 = -l_3\ .$$

Variant (30)

$$G = \begin{pmatrix} 0 & 2k_1 & n_0+n_3 & n_1-in_2 \\ 0 & 2k_0 & n_1+in_2 & n_0-n_3 \\ 0 & 2l_1 & 2m_0 & 2m_1 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$l_0 = 0, \qquad l_3 = 0, \qquad l_1 = -il_2\ ,$$

$$k_0 = -k_3, \qquad k_1 = -ik_2, \qquad m_1 = -im_2, \qquad m_0 = m_3\ .$$

Variant (31)

$$G = \begin{pmatrix} 2k_0 & 0 & n_0+n_3 & n_1-in_2 \\ 2k_1 & 0 & n_1+in_2 & n_0-n_3 \\ 2l_0 & 0 & 2m_0 & 2m_1 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$l_1 = 0, \qquad l_2 = 0, \qquad l_0 = l_3\ ,$$

$$k_0 = k_3, \qquad k_1 = ik_2, \qquad m_1 = -im_2, \qquad m_0 = m_3\ .$$

Variant (32)

$$G = \begin{pmatrix} k_0+k_3 & k_1-ik_2 & 0 & 2n_1 \\ k_1+ik_2 & k_0-k_3 & 0 & 2n_0 \\ 2l_0 & 2l_1 & 0 & 2m_1 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$m_0 = 0, \qquad m_3 = 0, \qquad m_1 = -im_2\ ,$$

$$l_0 = l_3, \qquad l_1 = -il_2, \qquad n_1 = -in_2, \qquad n_0 = -n_3\ .$$

Variant (33)

$$G = \begin{pmatrix} k_0+k_3 & k_1-ik_2 & 2n_0 & 0 \\ k_1+ik_2 & k_0-k_3 & 2n_1 & 0 \\ 2l_0 & 2l_1 & 2m_0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$m_1 = 0, \qquad m_2 = 0, \qquad m_0 = m_3\ ,$$

$$l_0 = l_3, \qquad l_1 = -il_2, \qquad n_1 = in_2, \qquad n_0 = n_3\ .$$

Let us remind the general form of the matrix G

$$G = \begin{pmatrix} k_0+k_3 & k_1-ik_2 & n_0+n_3 & n_1-in_2 \\ k_1+ik_2 & k_0-k_3 & n_1+in_2 & n_0-n_3 \\ l_0+l_3 & l_1-il_2 & m_0+m_3 & m_1-im_2 \\ l_1+il_2 & l_0-l_3 & m_1+im_2 & m_0-m_3 \end{pmatrix}.$$

15 Concluding Remarks

Let us summarize the results. The matrices of the form

$$G = \begin{pmatrix} k_0 + \mathbf{k}\,\vec{\sigma} & n_0 + \mathbf{n}\,\vec{\sigma} \\ l_0 + \mathbf{l}\,\vec{\sigma} & m_0 + \mathbf{m}\,\vec{\sigma} \end{pmatrix} ,$$

are described by their determinant [5]

$$\begin{aligned} \det\, G = (kk)\,(mm) + (nn)\,(ll) - 2\,(kn)\,(ml)\; - \\ 2\,(\,-k_0\,\mathbf{n} + n_0\,\mathbf{k} + i\,\mathbf{k}\times\mathbf{n}\,)\,(\,-m_0\,\mathbf{l} + l_0\,\mathbf{m} + i\,\mathbf{m}\times\mathbf{l}\,)\,; \end{aligned} \tag{15.1}$$

the classification of degenerate 4-dimensional matrices ($\det G = 0$), of rank 1, 2, 3 can be schematically described by

$$\begin{gathered} (k) \longrightarrow 7\,,\;\; (m) \longrightarrow 7\,,\quad (n) \longrightarrow 4\,,\quad (l) \longrightarrow 4\,, \\ (k,m) \longrightarrow 5\,,\qquad (l,n) \longrightarrow 2\,,\qquad (k,n) \longrightarrow 2\,, \\ (k,l) \longrightarrow 2\,,\qquad (n,m) \longrightarrow 2\,,\qquad (m,l) \longrightarrow 2\,, \\ (k,m,n) \longrightarrow 2\,,\qquad (k,m,l) \longrightarrow 2\,, \\ (n,l,k) \longrightarrow 2\,,\qquad (n,l,m) \longrightarrow 2\,. \end{gathered}$$

Thus, we produced many different subsets of real matrices, mainly with semi-group structure. These subset themselves can be of mathematical interest. The question of applying them as Mueller's type remains open.

Acknowledgment

The present work was developed under the auspices of Grant F14RA-006, within the cooperation framework between Romanian Academy and Belarusian Republican Foundation for Fundamental Research.

References

[1] Red'kov,V.M.; Bogush, A.A.; Tokarevskaya, N.G., On Parametrization of the Linear $GL(4,\mathbb{C})$ and unitary $SU(4)$ groups in terms of Dirac matrices. SIGMA. 2008. **4**, 021 [46 pages]; arXiv:0802.2634 [math-ph].

[2] Ovsiyuk, E.M.; Veko, O.V.; Red'kov, V.M., Mueller semi-groups of rank 1 and 2. Problems of Physics, Informatick, and Technics. 2012, no 2(11). 34–40 (in Russian).

[3] Ovsiyuk, E.M.; Red'kov, V.M., Is there possible the Finsler geometrization of the polarization optics? Hypecomplex Numbers in Geometry and Physics. 2012. **9**. no 1(17). 1–56 (in Russian); Report to: VIII-th International Conference Finsler Extensions of Relativity Theory (FERT-2012) June 25th-July 1st 2012 Moscow-Fryazino, Russia.

[4] Ovsiyuk, E.M.; Red'kov, V.M., Degenerate 4-dimensional matrices with semi-group structure and polarization optics. Vestnik RUDN. 2013. Ser. mathematics, physics. no 1. 245–259.

[5] Red'kov, V.M.; Bogush, A.A.; Tokarevskaya, N.G., (4×4)-matrices in Dirac parametrization: inversion problem and determinant. 32 pages, arXiv:0709.3574v2 [hep-th].